

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C12N 9/54</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/11770</b> <b>(43) International Publication Date:</b> 11 March 1999 (11.03.99)
<b>(21) International Application Number:</b> PCT/DK98/00361 <b>(22) International Filing Date:</b> 19 August 1998 (19.08.98) <b>(30) Priority Data:</b> 0988/97 29 August 1997 (29.08.97) DK <b>(71) Applicant (for all designated States except US):</b> NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> HANSEN, Peter, Kamp [DK/DK]; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsværd (DK). BAUDITZ, Peter [DK/DK]; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsværd (DK). MIKKELSEN, Frank [DK/DK]; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsværd (DK). ANDERSEN, Kim, Vilbour [DK/DK]; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsværd (DK). <b>(74) Common Representative:</b> NOVO NORDISK a/s; Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> PROTEASE VARIANTS AND COMPOSITIONS  <b>(57) Abstract</b>  Enzymes produced by mutating the genes for a number of subtilases and expressing the mutated genes in suitable hosts are presented. The enzymes exhibit improved wash performance in any detergent in comparison to their wild type parent enzymes.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Title: Protease Variants and Compositions

### TECHNICAL FIELD

5 This invention relates to novel mutant protease enzymes or enzyme variants useful in formulating detergent compositions and exhibiting improved wash performance in detergents; cleaning and detergent compositions containing said enzymes; mutated genes coding for the expression of said enzymes when  
10 inserted into a suitable host cell or organism; and such host cells transformed therewith and capable of expressing said enzyme variants.

### BACKGROUND OF THE INVENTION

15

In the detergent industry enzymes have for more than 30 years been implemented in washing formulations. Enzymes used in such formulations comprise proteases, lipases, amylases, cellulases, as well as other enzymes, or mixtures thereof. Commercially  
20 most important enzymes are proteases.

An increasing number of commercially used proteases are protein engineered variants of naturally occurring wild type proteases, e.g. DURAZYM<sup>®</sup> (Novo Nordisk A/S), RELEASE<sup>®</sup> (Novo Nordisk A/S), MAXAPEM<sup>®</sup> (Gist-Brocades N.V.), PURAFECT<sup>®</sup>  
25 (Genencor International, Inc.).

Further a number of protease variants are describe in the art, such as in EP 130756 (GENENTECH) (corresponding to US Reissue Patent No. 34,606 (GENENCOR)); EP 214435 (HENKEL); WO 87/04461 (AMGEN); WO 87/05050 (GENEX); EP 260105 (GENENCOR);  
30 Thomas, Russell, and Fersht (1985) *Nature* **318** 375-376; Thomas, Russell, and Fersht (1987) *J. Mol. Biol.* **193** 803-813; Russel and Fersht *Nature* **328** 496-500 (1987); WO 88/08028 (Genex); WO 88/08033 (Amgen); WO 95/27049 (SOLVAY S.A.); WO 95/30011 (PROCTER & GAMBLE COMPANY); WO 95/30010 (PROCTER & GAMBLE  
35 COMPANY); WO 95/29979 (PROCTER & GAMBLE COMPANY); US 5.543.302 (SOLVAY S.A.); EP 251 446 (GENENCOR); WO 89/06279 (NOVO NORDISK A/S); WO 91/00345 (NOVO NORDISK A/S); EP 525 610 A1 (SOLVAY);

WO 94/02618 (GIST-BROCADES N.V.); and WO 96/34946 (NOVO NORDISK A/S).

However, even though a number of useful protease variants have been described, there is still a need for new improved protease variants for a number of industrial uses.

Therefore, an object of the present invention, is to provide improved protein engineered protease variants, especially for use in the detergent industry.

10        SUMMARY OF THE INVENTION

The present inventors have intensively studied numerous of the possible combinations of the V95, G97 and A98 residues of SAVINASE®, and identified a number of variants with increased improved wash performance.

15        For further details reference is made to working examples herein (*vide infra*).

Accordingly, the present invention relates in its first aspect to a subtilase protease variant having improved wash performance in detergents, comprising modification(s) in position(s) 95, 97 and/or 98.

Preferably a subtilase variant according to the invention comprises modifications in both positions 97 and 98, and more preferred comprises modifications in all three positions 95, 97, and 98.

25        In a second aspect the invention relates to a subtilase enzyme variant having improved wash performance in detergents, comprising at least one modification chosen from the group comprising:

95T+97T+98E

30        97E+98V; or

a variant comprising one or more conservative modification(s) in any of the above mentioned variants (e.g. a conservative modification of a 97E(acidic a.a.)+98V variant include a variant such as 97D(acidic a.a.)+98V.).

35

In a third aspect the invention relates to an isolated DNA sequence encoding a subtilase variant of the invention.

In a fourth aspect the invention relates to an expression vector comprising an isolated DNA sequence encoding a subtilase variant of the invention.

In a fifth aspect the invention relates to a microbial  
5 host cell transformed with an expression vector according to the fourth aspect.

In a further aspect the invention relates to the production of the subtilisin enzymes of the invention by inserting an expression vector according to the fourth aspect  
10 into a suitable microbial host, cultivating the host to express the desired subtilase enzyme, and recovering the enzyme product.

Even further the invention relates to a composition comprising a subtilase variant of the invention.

15 Finally the invention relates to the use of the mutant enzymes for a number of industrial relevant uses, in particular for use in cleaning compositions and cleaning compositions comprising the mutant enzymes, especially detergent compositions comprising the mutant subtilisin enzymes.

20

#### DEFINITIONS

Prior to discussing this invention in further detail, the following term will first be defined.

Nomenclature of Amino Acids

	A	=	Ala	=	Alanine
	V	=	Val	=	Valine
5	L	=	Leu	=	Leucine
	I	=	Ile	=	Isoleucine
	P	=	Pro	=	Proline
	F	=	Phe	=	Phenylalanine
	W	=	Trp	=	Tryptophan
10	M	=	Met	=	Methionine
	G	=	Gly	=	Glycine
	S	=	Ser	=	Serine
	T	=	Thr	=	Threonine
	C	=	Cys	=	Cysteine
15	Y	=	Tyr	=	Tyrosine
	N	=	Asn	=	Asparagine
	Q	=	Gln	=	Glutamine
	D	=	Asp	=	Aspartic Acid
	E	=	Glu	=	Glutamic Acid
20	K	=	Lys	=	Lysine
	R	=	Arg	=	Arginine
	H	=	His	=	Histidine
	X	=	Xaa	=	Any amino acid

Nomenclature of nucleic acids

25	A	=	Adenine
	G	=	Guanine
	C	=	Cytosine
	T	=	Thymine (only in DNA)
	U	=	Uracil (only in RNA)

30

Nomenclature of variants

In describing the various enzyme variants produced or contemplated according to the invention, the following nomenclatures have been adapted for ease of reference:

35

Original amino acid(s) position(s) substituted amino acid(s)

According to this the substitution of Glutamic acid for glycine in position 195 is designated as:

Gly 195 Glu or G195E

a deletion of glycine in the same position is:

5 Gly 195 \* or G195\*

and insertion of an additional amino acid residue such as lysine is:

Gly 195 GlyLys or G195GK

10 Where a deletion in comparison with the sequence used for the numbering is indicated, an insertion in such a position is indicated as:

\* 36 Asp or \*36D

for insertion of an aspartic acid in position 36

15

Multiple mutations are separated by pluses, i.e.:

Arg 170 Tyr + Gly 195 Glu or R170Y+G195E

representing mutations in positions 170 and 195 substituting tyrosine and glutamic acid for arginine and glycine, respectively.  
20

### Proteases

Enzymes cleaving the amide linkages in protein substrates are classified as proteases, or (interchangeably) peptidases (see  
25 Walsh, 1979, *Enzymatic Reaction Mechanisms*. W.H. Freeman and Company, San Francisco, Chapter 3).

### Numbering of amino acid positions/residues

If no other mentioned the amino acid numbering used herein  
30 correspond to that of the subtilase BPN<sup>-</sup> (BASBPN) sequence. For further description of the BPN<sup>-</sup> sequence see Siezen et al., *Protein Engng.* 4 (1991) 719-737 and Figure 1.

### Serine proteases

35 A serine protease is an enzyme which catalyzes the hydrolysis of peptide bonds, and in which there is an essential serine residue at the active site (White, Handler and Smith, 1973

"Principles of Biochemistry," Fifth Edition, McGraw-Hill Book Company, NY, pp. 271-272).

The bacterial serine proteases have molecular weights in the 20,000 to 45,000 Daltons range. They are inhibited by diisopropylfluorophosphate. They hydrolyze simple terminal esters and are similar in activity to eukaryotic chymotrypsin, also a serine protease. A more narrow term, alkaline protease, covering a sub-group, reflects the high pH optimum of some of the serine proteases, from pH 9.0 to 11.0 (for review, see Priest (1977) *Bacteriological Rev.* 41 711-753).

#### Subtilases

A sub-group of the serine proteases tentatively designated subtilases has been proposed by Siezen et al., *Protein Engng.* 4 (1991) 719-737. They are defined by homology analysis of more than 40 amino acid sequences of serine proteases previously referred to as subtilisin-like proteases. A subtilisin was previously defined as a serine protease produced by Gram-positive bacteria or fungi, and according to Siezen et al. now is a subgroup of the subtilases. A wide variety of subtilases have been identified, and the amino acid sequence of a number of subtilases have been determined. For a more detailed description of such subtilases and their amino acid sequences reference is made to Siezen et al. and figure 1 herein.

One subgroup of the subtilases, I-S1, comprises the "classical" subtilisins, such as subtilisin 168, subtilisin BPN', subtilisin Carlsberg (ALCALASE®, NOVO NORDISK A/S), and subtilisin DY.

A further subgroup of the subtilases I-S2, is recognised by Siezen et al. (*supra*). Sub-group I-S2 proteases are described as highly alkaline subtilisins and comprise enzymes such as subtilisin PB92 (MAXACAL®, Gist-Brocades NV), subtilisin 309 (SAVINASE®, NOVO NORDISK A/S), subtilisin 147 (ESPERASE®, NOVO NORDISK A/S), and alkaline elastase YaB.

35

#### "SAVINASE®"

SAVINASE® is marketed by NOVO NORDISK A/S.



It is subtilisin 309 from B. Lentus and differs from BABP92 only in having N87S (see figure 1 herein).

#### Parent subtilase

5 The term "parent subtilase" is a subtilase defined according to Siezen et al. (Protein Engineering 4:719-737 (1991)). For further details see description of "SUBTILASES" immediately above. A parent subtilase may also be a subtilase isolated from a natural source, wherein subsequent modification have been made  
10 while retaining the characteristic of a subtilase. Alternatively the term "parent subtilase" may be termed "wild-type subtilase".

#### Modification(s) of a subtilase variant

15 The term "modification(s)" used in connection with modification(s) of a subtilase variant as discussed herein is defined to include chemical modification as well as genetic manipulation. The modification(s) can be by substitution, deletion and/or insertions in or at the amino acid(s) of interest.

20

#### Subtilase variant

In the context of this invention, the term subtilase variant or mutated subtilase means a subtilase that has been produced by an organism which is expressing a mutant gene derived from a  
25 parent microorganism which possessed an original or parent gene and which produced a corresponding parent enzyme, the parent gene having been mutated in order to produce the mutant gene from which said mutated subtilase protease is produced when expressed in a suitable host.

30

#### Homologous subtilase sequences

Specific amino acid residues of SAVINASE® subtilase are identified for modification herein to obtain a subtilase variant of the invention.

35

However, the invention is not limited to modifications of this particular subtilase, but extend to other parent (wild-

type) subtilases, which have a homologous primary structure to that of SAVINASE®.

In order to identify other homologous subtilases, within the scope of this invention, an alignment of said 5 subtilase(s) to a group of previously aligned subtilases is performed keeping the previous alignment constant. A comparison to 18 highly conserved residues in subtilases is performed. The 18 highly conserved residues are shown in table I (see Siezen et al. for further details relating to said conserved 10 residues).

Table I

18 highly conserved residues in subtilases

	Position:	Conserved residue
15	23	G
	32	D
	34	G
	39	H
	64	H
20	65	G
	66	T
	70	G
	83	G
	125	S
25	127	G
	146	G
	154	G
	155	N
	219	G
30	220	T
	221	S
	225	P

After aligning allowing for necessary insertions and 35 deletions in order to maintain the alignment suitable homologous residues are identified. Said homologous residues can then be modified according to the invention.

Using the CLUSTALW (version 1.5, April 1995) computer alignment program (Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994) Nucleic Acids Research, 22:4673-4680.), with GAP open penalty of 10.0 and GAP extension penalty of 0.1, using  
 5 the BLOSUM30 protein weight matrix, alignment of a given subtilase to a group of previously aligned subtilases is achieved using the *Profile alignments* option in the program. For a given subtilase to be within the scope of the invention, preferably 100% of the 18 highly conserved residues should be  
 10 conserved. However, alignment of greater than or equal to 17 out of the 18 residues, or as little as 16 of said conserved residues is also adequate to identify homologous residues. Conservation of the, in subtilases, catalytic triad Asp32/His64/Ser221 should be maintained.

15 The previously defined alignment is shown figure 1, where the percent identity of the individual subtilases in this alignment to the 18 highly conserved residues are shown too.

Based on this description it is routine for a person skilled in the art to identify suitable homologous subtilases  
 20 and corresponding homologous residues, which can be modified according to the invention. To illustrate this table II below shows a limited list a homologous subtilases and corresponding suitable residues to be modified according to the invention.

25 Table II

Homologous Subtilases and corresponding homologous residues, suitable to be modified according to the invention.

Pos\Enz.	BASBPN	BYSYAB	BLS309	BLS147	TVTHER
97+98	G97E+	G97E+	G97E+	D97E+	D97E+
	A98V	A98V	A98V	R98V	N98V
95+97+98	V95T+	V95T+	V95T+	V95T+	V95T+
	G97T+	G97T+	G97T+	D97T+	D97T+
	A98E	A98E	A98E	R98E	N98E

It is obvious that a similar or larger table covering other homologous subtilases may easily be produced by a person skilled in the art.

5 Wash performance

The ability of an enzyme to catalyze the degradation of various naturally occurring substrates present on the objects to be cleaned during e.g. wash is often referred to as its washing ability, washability, detergency, or wash performance.  
10 Throughout this application the term wash performance will be used to encompass this property.

Isolated DNA sequence

The term "isolated", when applied to a DNA sequence  
15 molecule, denotes that the DNA sequence has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated  
20 from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example, Dynan and Tijan, Nature 316:774-78, 1985). The term "an isolated DNA sequence" may alternatively be termed "a cloned DNA sequence".  
25

30 Isolated protein

When applied to a protein, the term "isolated" indicates that the protein is found in a condition other than its native environment. In a preferred form, the isolated protein is substantially free of other proteins, particularly  
35 other homologous proteins (i.e. "homologous impurities" (see below)). It is preferred to provide the protein in a highly purified form, i.e., greater than 40% pure, greater than 60% pure, greater than 80% pure, more preferably greater than 95%

pure, and even more preferably greater than 99% pure, as determined by SDS-PAGE. The term "isolated protein" may alternatively be termed "purified protein".

5 Homologous impurities

The term "homologous impurities" means any impurity (e.g. another polypeptide than the polypeptide of the invention) which originate from the homologous cell where the polypeptide of the invention is originally obtained from.

10

Obtained from

The term "obtained from" as used herein in connection with a specific microbial source, means that the polynucleotide and/or polypeptide produced by the specific source, or by a cell in  
15 which a gene from the source have been inserted.

Substrate

The term "Substrate" used in connection with a substrate for a protease is should be interpreted in its broadest form as  
20 comprising a compound containing at least one peptide bond susceptible to hydrolysis by a subtilisin protease.

Product

The term "product" used in connection with a product derived  
25 from a protease enzymatic reaction should in the context of this invention be interpreted to include the products of a hydrolysis reaction involving a subtilase protease. A product may be the substrate in a subsequent hydrolysis reaction.

30

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows an alignment of a number of homologous subtilases, which are aligned to 18 highly conserved residues in subtilases. 18 highly conserved residues are highlighted in bold. All shown  
35 subtilases, except JP170, have 100% identity in said conserved residues. JP170 is having an "N" in stead of "G" in conserved residues G146.

**DETAILED DESCRIPTION OF THE INVENTION****Subtilase variants with improved wash performance:**

5           The present inventors have identified the improved wash performance variants in BLS309 (SAVINASE®).

          Accordingly, an embodiment of the invention relates to a subtilase enzyme variant, wherein the modification is chosen from the group comprising:

10           V95T+G97T+A98E  
          G97E+A98V; or

a variant comprising one or more conservative modification(s) in any of the above mentioned variants (e.g. a conservative modification of a G97E(acidic a.a.)+A98V variant include a  
15 variant such as G97D(acidic a.a.)+A98V.).

Numerous subtilase variants of the invention is tested herein and showing improved wash-performance in detergents (see working examples herein (*vide infra*)).

It is well known in the art that substitution of one  
20 amino acid to a similar conservative amino acid only give a minor change in the characteristic of the enzyme.

Table III below list groups of conservative amino acids.

25

**Table III****Conservative amino acid substitutions**

	Basic:	arginine
		lysine
		histidine
30	Acidic:	glutamic acid
		aspartic acid
	Polar:	glutamine
		asparagine
35	Hydrophobic:	leucine
		isoleucine
		valine
	Aromatic:	phenylalanine

tryptophan  
tyrosine  
Small: glycine  
alanine  
5 serine  
threonine  
methionine

Accordingly, subtilase variants such as 97E+98V and  
10 97D+98V will have a similar wash-performance improvement.

Further, subtilase variants such as G97E+A98V and  
G97D+A98V will have a similar wash-performance improvement too.

Based on the disclosed subtilase variants herein, it is  
routine work, for a person skilled in the art, to identify  
15 further suitable conservative substitutions in order to obtain  
a subtilase variant with improved wash-performance.

In embodiments of the invention, the subtilases of  
interest are those belonging to the subgroups I-S1 and I-S2.

Relating to subgroup I-S1 preferred parent subtilase is  
20 chosen from the group comprising ABSS168, BASBPN, BSSDY, and  
BLSCAR or functional variants thereof having retained the  
characteristic of sub-group I-S1.

Relating to subgroup I-S2 preferred parent subtilase is  
chosen from the group comprising BLS147, BLS309, BAPB92, TVTHER  
25 AND BYSYAB or functional variants thereof having retained the  
characteristic of sub-group I-S2.

The present invention also comprises any one or more  
modifications in the above mentioned positions in combination  
with any other modification to the amino acid sequence of the  
30 parent enzyme. Especially combinations with other modifications  
known in the art to provide improved properties to the enzyme  
are envisaged. The art describe a number of subtilase variants  
with different improved properties and a number of those are  
mentioned in the "Background of the invention" section herein  
35 (*vide supra*). Those references are disclosed here as references  
to identify a subtilase variant, which advantageously can be  
combined with a subtilase variant of the invention.

Such combinations comprise the positions: 222 (improve oxidation stability), 218 (improves thermal stability), substitutions in the Ca-binding sites stabilising the enzyme, e.g. position 76, and many other apparent from the prior art.

5 In further embodiments a subtilase variant of the invention may advantageously be combined with one or more modification(s) in any of the positions:

27, 36, 57, 76, 97, 101, 104, 120, 123, 167, 170, 206, 218, 222, 224, 235 and 274.

10 Specifically the following BLS309 and BAPB92 variants are considered appropriate for combination:

K27R, \*36D, S57P, N76D, G97N, S101G, V104A, V104N, V104Y, H120D, N123S, Y167A, Y167I, R170S, R170L, R170N, Q206E, N218S, M222S, M222A, T224S, K235L and T274A.

15 Furthermore variants comprising any of the variants V104N+S101G, K27R+V104Y+N123S+T274A, or N76D+V104A or other combinations of these mutations (V104N, S101G, K27R, V104Y, N123S, T274A, N76D, V104A), in combination with any one or more of the modification(s) mentioned above exhibit improved  
20 properties.

Even further subtilase variants of the main aspect(s) of the invention are preferably combined with one or more modification(s) in any of the positions 129, 131, 133 and 194, preferably as 129K, 131H, 133P, 133D and 194P modifications,  
25 and most preferably as P129K, P131H, A133P, A133D and A194P modifications. Any of those modification(s) may give a higher expression level of a subtilase variant of the invention.

#### PRODUCING MUTATIONS IN SUBTILASE GENES

30 Many methods for cloning a subtilase of the invention and for introducing mutations into genes (e.g. subtilase genes) are well known in the art.

In general standard procedures for cloning of genes and introducing mutations (random and/or site directed) into said  
35 genes may be used in order to obtain a subtilase variant of the invention. For further description of suitable techniques reference is made to working examples herein (*vide infra*) and (Sambrook et al. (1989) Molecular cloning: A laboratory manual,



Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995; Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for Bacillus". John Wiley and  
5 Sons, 1990); and WO 96/34946.

#### EXPRESSION VECTORS

A recombinant expression vector comprising a DNA construct encoding the enzyme of the invention may be any  
10 vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which  
15 is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome in part or in its entirety and replicated together with the chromosome(s) into which it has been integrated.

20 The vector is preferably an expression vector in which the DNA sequence encoding the enzyme of the invention is operably linked to additional segments required for transcription of the DNA. In general, the expression vector is derived from plasmid or viral DNA, or may contain elements of  
25 both. The term, "operably linked" indicates that the segments are arranged so that they function in concert for their intended purposes, e.g. transcription initiates in a promoter and proceeds through the DNA sequence coding for the enzyme.

The promoter may be any DNA sequence which shows  
30 transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell.

Examples of suitable promoters for use in bacterial host cells include the promoter of the *Bacillus*  
35 *stearothermophilus* maltogenic amylase gene, the *Bacillus licheniformis* alpha-amylase gene, the *Bacillus amyloliquefaciens* alpha-amylase gene, the *Bacillus subtilis* alkaline protease gen, or the *Bacillus pumilus* xylosidase gene,

or the phage Lambda P<sub>R</sub> or P<sub>L</sub> promoters or the E. coli lac, trp or tac promoters.

The DNA sequence encoding the enzyme of the invention may also, if necessary, be operably connected to a suitable  
5 terminator.

The recombinant vector of the invention may further comprise a DNA sequence enabling the vector to replicate in the host cell in question.

The vector may also comprise a selectable marker, e.g.  
10 a gene the product of which complements a defect in the host cell, or a gene encoding resistance to e.g. antibiotics like kanamycin, chloramphenicol, erythromycin, tetracycline, spectinomycine, or the like, or resistance to heavy metals or herbicides.

15 To direct an enzyme of the present invention into the secretory pathway of the host cells, a secretory signal sequence (also known as a leader sequence, prepro sequence or pre sequence) may be provided in the recombinant vector. The secretory signal sequence is joined to the DNA sequence  
20 encoding the enzyme in the correct reading frame. Secretory signal sequences are commonly positioned 5' to the DNA sequence encoding the enzyme. The secretory signal sequence may be that normally associated with the enzyme or may be from a gene encoding another secreted protein.

25 The procedures used to ligate the DNA sequences coding for the present enzyme, the promoter and optionally the terminator and/or secretory signal sequence, respectively, or to assemble these sequences by suitable PCR amplification schemes, and to insert them into suitable vectors containing the  
30 information necessary for replication or integration, are well known to persons skilled in the art (cf., for instance, Sambrook et al., op.cit.).

#### HOST CELL

35 The DNA sequence encoding the present enzyme introduced into the host cell may be either homologous or heterologous to the host in question. If homologous to the host cell, i.e. produced by the host cell in nature, it will typically be operably

connected to another promoter sequence or, if applicable, another secretory signal sequence and/or terminator sequence than in its natural environment. The term "homologous" is intended to include a DNA sequence encoding an enzyme native to the host organism in question. The term "heterologous" is intended to include a DNA sequence not expressed by the host cell in nature. Thus, the DNA sequence may be from another organism, or it may be a synthetic sequence.

The host cell into which the DNA construct or the recombinant vector of the invention is introduced may be any cell which is capable of producing the present enzyme and includes bacteria, yeast, fungi and higher eukaryotic cells.

Examples of bacterial host cells which, on cultivation, are capable of producing the enzyme of the invention are gram-positive bacteria such as strains of *Bacillus*, such as strains of *B. subtilis*, *B. licheniformis*, *B. lentus*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. amyloliquefaciens*, *B. coagulans*, *B. circulans*, *B. lautus*, *B. megatherium* or *B. thuringiensis*, or strains of *Streptomyces*, such as *S. lividans* or *S. murinus*, or gram-negative bacteria such as *Escherichia coli*. The transformation of the bacteria may be effected by protoplast transformation, electroporation, conjugation, or by using competent cells in a manner known per se (cf. Sambrook et al., supra).

When expressing the enzyme in bacteria such as *E. coli*, the enzyme may be retained in the cytoplasm, typically as insoluble granules (known as inclusion bodies), or may be directed to the periplasmic space by a bacterial secretion sequence. In the former case, the cells are lysed and the granules are recovered and denatured after which the enzyme is refolded by diluting the denaturing agent. In the latter case, the enzyme may be recovered from the periplasmic space by disrupting the cells, e.g. by sonication or osmotic shock, to release the contents of the periplasmic space and recovering the enzyme.

When expressing the enzyme in gram-positive bacteria such as *Bacillus* or *Streptomyces* strains, the enzyme may be retained in the cytoplasm, or may be directed to the

extracellular medium by a bacterial secretion sequence. In the latter case, the enzyme may be recovered from the medium as described below.

#### 5 METHOD OF PRODUCING SUBTILASE

The present invention provides a method of producing an isolated enzyme according to the invention, wherein a suitable host cell, which has been transformed with a DNA sequence encoding the enzyme, is cultured under conditions permitting  
10 the production of the enzyme, and the resulting enzyme is recovered from the culture.

When an expression vector comprising a DNA sequence encoding the enzyme is transformed into a heterologous host cell it is possible to enable heterologous recombinant  
15 production of the enzyme of the invention.

Thereby it is possible to make a highly purified subtilase composition, characterized in being free from homologous impurities.

In this context homologous impurities means any  
20 impurities (e.g. other polypeptides than the enzyme of the invention) which originate from the homologous cell where the enzyme of the invention is originally obtained from.

The medium used to culture the transformed host cells may be any conventional medium suitable for growing the host  
25 cells in question. The expressed subtilase may conveniently be secreted into the culture medium and may be recovered therefrom by well-known procedures including separating the cells from the medium by centrifugation or filtration, precipitating proteinaceous components of the medium by means of a salt such  
30 as ammonium sulphate, followed by chromatographic procedures such as ion exchange chromatography, affinity chromatography, or the like.

#### USE OF A SUBTILASE VARIANT OF THE INVENTION

35 A subtilase protease variant of the invention may be used for a number of industrial applications, in particular within the detergent industry.

Further the invention relates to an enzyme composition, which comprise a subtilase variant of the invention.

An summary of preferred industrial applications and corresponding preferred enzyme compositions are described  
5 below.

This summary is not in any way intended to be a complete list of suitable applications of a subtilase variant of the invention. A subtilase variants of the invention may be used in other industrial applications known in the art to include use  
10 of a protease, in particular a subtilase.

#### DETERGENT COMPOSITIONS COMPRISING THE MUTANT ENZYMES

The present invention comprises the use of the mutant enzymes of the invention in cleaning and detergent compositions  
15 and such compositions comprising the mutant subtilisin enzymes. Such cleaning and detergent compositions are well described in the art and reference is made to WO 96/34946; WO 97/07202; WO 95/30011 for further description of suitable cleaning and detergent compositions.

20 Further reference is made to workings example(s) herein showing wash performance improvements for a number of subtilase variants of the invention.

#### DETERGENT DISCLOSURE AND EXAMPLES

##### 25 Surfactant system

The detergent compositions according to the present invention comprise a surfactant system, wherein the surfactant can be selected from nonionic and/or anionic and/or cationic and/or ampholytic and/or zwitterionic and/or semi-polar  
30 surfactants.

The surfactant is typically present at a level from 0.1% to 60% by weight.

The surfactant is preferably formulated to be compatible with enzyme components present in the composition. In liquid or  
35 gel compositions the surfactant is most preferably formulated in such a way that it promotes, or at least does not degrade, the stability of any enzyme in these compositions.

Preferred systems to be used according to the present invention comprise as a surfactant one or more of the nonionic and/or anionic surfactants described herein.


Polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols are suitable for use as the nonionic surfactant of the surfactant systems of the present invention, with the polyethylene oxide condensates being preferred. These compounds include the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 14 carbon atoms, preferably from about 8 to about 14 carbon atoms, in either a straight chain or branched-chain configuration with the alkylene oxide. In a preferred embodiment, the ethylene oxide is present in an amount equal to from about 2 to about 25 moles, more preferably from about 3 to about 15 moles, of ethylene oxide per mole of alkyl phenol. Commercially available nonionic surfactants of this type include Igepal<sup>TM</sup> CO-630, marketed by the GAF Corporation; and Triton<sup>TM</sup> X-45, X-114, X-100 and X-102, all marketed by the Rohm & Haas Company. These surfactants are commonly referred to as alkylphenol alkoxylates (e.g., alkyl phenol ethoxylates).

The condensation products of primary and secondary aliphatic alcohols with about 1 to about 25 moles of ethylene oxide are suitable for use as the nonionic surfactant of the nonionic surfactant systems of the present invention. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from about 8 to about 22 carbon atoms. Preferred are the condensation products of alcohols having an alkyl group containing from about 8 to about 20 carbon atoms, more preferably from about 10 to about 18 carbon atoms, with from about 2 to about 10 moles of ethylene oxide per mole of alcohol. About 2 to about 7 moles of ethylene oxide and most preferably from 2 to 5 moles of ethylene oxide per mole of alcohol are present in said condensation products. Examples of commercially available nonionic surfactants of this type include Tergitol<sup>TM</sup> 15-S-9 (The condensation product of C<sub>11</sub>-C<sub>15</sub> linear alcohol with 9 moles ethylene oxide), Tergitol<sup>TM</sup> 24-L-6 NMW (the condensation product of C<sub>12</sub>-C<sub>14</sub> primary alcohol with 6

moles ethylene oxide with a narrow molecular weight distribution), both marketed by Union Carbide Corporation; Neodol<sup>TM</sup> 45-9 (the condensation product of C<sub>14</sub>-C<sub>15</sub> linear alcohol with 9 moles of ethylene oxide), Neodol<sup>TM</sup> 23-3 (the  
5 condensation product of C<sub>12</sub>-C<sub>13</sub> linear alcohol with 3.0 moles of ethylene oxide), Neodol<sup>TM</sup> 45-7 (the condensation product of C<sub>14</sub>-C<sub>15</sub> linear alcohol with 7 moles of ethylene oxide), Neodol<sup>TM</sup> 45-5 (the condensation product of C<sub>14</sub>-C<sub>15</sub> linear alcohol with 5 moles of ethylene oxide) marketed by Shell Chemical Company,  
10 Kyro<sup>TM</sup> EOB (the condensation product of C<sub>13</sub>-C<sub>15</sub> alcohol with 9 moles ethylene oxide), marketed by The Procter & Gamble Company, and Genapol LA 050 (the condensation product of C<sub>12</sub>-C<sub>14</sub> alcohol with 5 moles of ethylene oxide) marketed by Hoechst. Preferred range of HLB in these products is from 8-11 and most  
15 preferred from 8-10.

Also useful as the nonionic surfactant of the surfactant systems of the present invention are alkylpolysaccharides disclosed in US 4,565,647, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably  
20 from about 10 to about 16 carbon atoms and a polysaccharide, e.g. a polyglycoside, hydrophilic group containing from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used,  
25 e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties (optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside). The intersaccharide bonds can be, e.g., between  
30 the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6- positions on the preceding saccharide units.

The preferred alkylpolyglycosides have the formula



35

wherein R<sup>2</sup> is selected from the group consisting of alkyl, alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which the alkyl groups contain from about 10 to

about 18, preferably from about 12 to about 14, carbon atoms; n is 2 or 3, preferably 2; t is from 0 to about 10, preferably 0; and x is from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7.

5 The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted with glucose, or a source of glucose, to form the glucoside (attachment at the 1-position). The additional glycosyl units can then be attached between  
10 their 1-position and the preceding glycosyl units 2-, 3-, 4-, and/or 6-position, preferably predominantly the 2-position.

The condensation products of ethylene oxide with a hydrophobic base formed by the condensation of propylene oxide with propylene glycol are also suitable for use as the  
15 additional nonionic surfactant systems of the present invention. The hydrophobic portion of these compounds will preferably have a molecular weight from about 1500 to about 1800 and will exhibit water insolubility. The addition of polyoxyethylene moieties to this hydrophobic portion tends to  
20 increase the water solubility of the molecule as a whole, and the liquid character of the product is retained up to the point where the polyoxyethylene content is about 50% of the total weight of the condensation product, which corresponds to condensation with up to about 40 moles of ethylene oxide.  
25 Examples of compounds of this type include certain of the commercially available Pluronic<sup>TM</sup> surfactants, marketed by BASF.

Also suitable for use as the nonionic surfactant of the nonionic surfactant system of the present invention, are the  
30 condensation products of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylenediamine. The hydrophobic moiety of these products consists of the reaction product of ethylenediamine and excess propylene oxide, and generally has a molecular weight of from  
35 about 2500 to about 3000. This hydrophobic moiety is condensed with ethylene oxide to the extent that the condensation product contains from about 40% to about 80% by weight of polyoxyethylene and has a molecular weight of from about 5,000




to about 11,000. Examples of this type of nonionic surfactant include certain of the commercially available Tetronic<sup>TM</sup> compounds, marketed by BASF.

Preferred for use as the nonionic surfactant of the surfactant systems of the present invention are polyethylene oxide condensates of alkyl phenols, condensation products of primary and secondary aliphatic alcohols with from about 1 to about 25 moles of ethyleneoxide, alkylpolysaccharides, and mixtures hereof. Most preferred are C<sub>8</sub>-C<sub>14</sub> alkyl phenol ethoxylates having from 3 to 15 ethoxy groups and C<sub>8</sub>-C<sub>18</sub> alcohol ethoxylates (preferably C<sub>10</sub> avg.) having from 2 to 10 ethoxy groups, and mixtures thereof.

Highly preferred nonionic surfactants are polyhydroxy fatty acid amide surfactants of the formula

15



wherein R<sup>1</sup> is H, or R<sup>1</sup> is C<sub>1-4</sub> hydrocarbyl, 2-hydroxyethyl, 2-hydroxypropyl or a mixture thereof, R<sup>2</sup> is C<sub>5-31</sub> hydrocarbyl, and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxylated derivative thereof. Preferably, R<sup>1</sup> is methyl, R<sup>2</sup> is straight C<sub>11-15</sub> alkyl or C<sub>16-18</sub> alkyl or alkenyl chain such as coconut alkyl or mixtures thereof, and Z is derived from a reducing sugar such as glucose, fructose, maltose or lactose, in a reductive amination reaction.

Highly preferred anionic surfactants include alkyl alkoxylated sulfate surfactants. Examples hereof are water soluble salts or acids of the formula RO(A)<sub>m</sub>SO<sub>3</sub>M wherein R is an unsubstituted C<sub>10</sub>-C<sub>24</sub> alkyl or hydroxyalkyl group having a C<sub>10</sub>-C<sub>24</sub> alkyl component, preferably a C<sub>12</sub>-C<sub>20</sub> alkyl or hydroxyalkyl, more preferably C<sub>12</sub>-C<sub>18</sub> alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a cation which can be, for

example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxylated sulfates as well as alkyl propoxylated sulfates are contemplated herein. Specific

5 examples of substituted ammonium cations include methyl-, dimethyl, trimethyl-ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and those derived from alkylamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the like.

10 Exemplary surfactants are C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (1.0) sulfate (C<sub>12</sub>-C<sub>18</sub>E(1.0)M), C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (2.25) sulfate (C<sub>12</sub>-C<sub>18</sub>(2.25)M, and C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (3.0) sulfate (C<sub>12</sub>-C<sub>18</sub>E(3.0)M), and C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (4.0) sulfate (C<sub>12</sub>-C<sub>18</sub>E(4.0)M), wherein M is conveniently selected


15 from sodium and potassium.

Suitable anionic surfactants to be used are alkyl ester sulfonate surfactants including linear esters of C<sub>8</sub>-C<sub>20</sub> carboxylic acids (i.e., fatty acids) which are sulfonated with gaseous SO<sub>3</sub> according to "The Journal of the American Oil

20 Chemists Society", 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from tallow, palm oil, etc.

The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprise alkyl ester

25 sulfonate surfactants of the structural formula:



wherein R<sup>3</sup> is a C<sub>8</sub>-C<sub>20</sub> hydrocarbyl, preferably an alkyl, or combination thereof, R<sup>4</sup> is a C<sub>1</sub>-C<sub>6</sub> hydrocarbyl, preferably an

35 alkyl, or combination thereof, and M is a cation which forms a water soluble salt with the alkyl ester sulfonate. Suitable salt-forming cations include metals such as sodium, potassium,

and lithium, and substituted or unsubstituted ammonium cations, such as monoethanolamine, diethanolamine, and triethanolamine. Preferably,  $R^3$  is  $C_{10}$ - $C_{16}$  alkyl, and  $R^4$  is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates  
5 wherein  $R^3$  is  $C_{10}$ - $C_{16}$  alkyl.

Other suitable anionic surfactants include the alkyl sulfate surfactants which are water soluble salts or acids of the formula  $ROSO_3M$  wherein R preferably is a  $C_{10}$ - $C_{24}$  hydrocarbyl, preferably an alkyl or hydroxyalkyl having a  $C_{10}$ -  
10  $C_{20}$  alkyl component, more preferably a  $C_{12}$ - $C_{18}$  alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g. sodium, potassium, lithium), or ammonium or substituted ammonium (e.g. methyl-, dimethyl-, and trimethyl ammonium cations and quaternary ammonium cations such as  
15 tetramethyl-ammonium and dimethyl piperdinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like). Typically, alkyl chains of  $C_{12}$ - $C_{16}$  are preferred for lower wash temperatures (e.g. below about  $50^\circ C$ ) and  $C_{16}$ - $C_{18}$   
20 alkyl chains are preferred for higher wash temperatures (e.g. above about  $50^\circ C$ ).

Other anionic surfactants useful for deterative purposes can also be included in the laundry detergent compositions of the present invention. These can include salts (including, for  
25 example, sodium, potassium, ammonium, and substituted ammonium salts such as mono- di- and triethanolamine salts) of soap,  $C_8$ - $C_{22}$  primary or secondary alkanesulfonates,  $C_8$ - $C_{24}$  olefinsulfonates, sulfonated polycarboxylic acids prepared by sulfonation of the pyrolyzed product of alkaline earth metal  
30 citrates, e.g., as described in British patent specification No. 1,082,179,  $C_8$ - $C_{24}$  alkylpolyglycoethersulfates (containing up to 10 moles of ethylene oxide); alkyl glycerol sulfonates, fatty acyl glycerol sulfonates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin  
35 sulfonates, alkyl phosphates, isethionates such as the acyl isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinates (especially saturated and unsaturated  $C_{12}$ - $C_{18}$  monoesters) and diesters of

sulfosuccinates (especially saturated and unsaturated C<sub>6</sub>-C<sub>12</sub> diesters), acyl sarcosinates, sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), branched primary  
 5 alkyl sulfates, and alkyl polyethoxy carboxylates such as those of the formula RO(CH<sub>2</sub>CH<sub>2</sub>O)<sub>k</sub>-CH<sub>2</sub>COO-M<sup>+</sup> wherein R is a C<sub>8</sub>-C<sub>22</sub> alkyl, k is an integer from 1 to 10, and M is a soluble salt forming cation. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin  
 10 acids and hydrogenated resin acids present in or derived from tall oil.

Alkylbenzene sulfonates are highly preferred. Especially preferred are linear (straight-chain) alkyl benzene sulfonates (LAS) wherein the alkyl group preferably contains from 10 to 18  
 15 carbon atoms.


Further examples are described in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in US 3,929,678, (Column 23, line 58 through Column 29, line 23,  
 20 herein incorporated by reference).

When included therein, the laundry detergent compositions of the present invention typically comprise from about 1% to about 40%, preferably from about 3% to about 20% by weight of such anionic surfactants.

25 The laundry detergent compositions of the present invention may also contain cationic, ampholytic, zwitterionic, and semi-polar surfactants, as well as the nonionic and/or anionic surfactants other than those already described herein.


Cationic deterative surfactants suitable for use in the  
 30 laundry detergent compositions of the present invention are those having one long-chain hydrocarbyl group. Examples of such cationic surfactants include the ammonium surfactants such as alkyltrimethylammonium halogenides, and those surfactants having the formula:

35



wherein  $R^2$  is an alkyl or alkyl benzyl group having from about 8 to about 18 carbon atoms in the alkyl chain, each  $R^3$  is selected from the group consisting of  $-\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}(\text{CH}_3)-$ ,  $-\text{CH}_2\text{CH}(\text{CH}_2\text{OH})-$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ , and mixtures thereof; each  $R^4$  is  
 5 selected from the group consisting of  $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_1$ - $\text{C}_4$  hydroxyalkyl, benzyl ring structures formed by joining the two  $R^4$  groups,  $-\text{CH}_2\text{CHOHCHOHCOR}^6\text{CHOHCH}_2\text{OH}$ , wherein  $R^6$  is any hexose or hexose polymer having a molecular weight less than about 1000, and hydrogen when  $y$  is not 0;  $R^5$  is the same as  $R^4$  or is  
 10 an alkyl chain, wherein the total number of carbon atoms or  $R^2$  plus  $R^5$  is not more than about 18; each  $y$  is from 0 to about 10, and the sum of the  $y$  values is from 0 to about 15; and  $X$  is any compatible anion.

Highly preferred cationic surfactants are the water  
 15 soluble quaternary ammonium compounds useful in the present composition having the formula:



20 wherein  $\text{R}_1$  is  $\text{C}_8$ - $\text{C}_{16}$  alkyl, each of  $\text{R}_2$ ,  $\text{R}_3$  and  $\text{R}_4$  is independently  $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_1$ - $\text{C}_4$  hydroxy alkyl, benzyl, and  $-(\text{C}_2\text{H}_{40})_x\text{H}$  where  $x$  has a value from 2 to 5, and  $X$  is an anion. Not more than one of  $\text{R}_2$ ,  $\text{R}_3$  or  $\text{R}_4$  should be benzyl.

The preferred alkyl chain length for  $\text{R}_1$  is  $\text{C}_{12}$ - $\text{C}_{15}$ ,  
 25 particularly where the alkyl group is a mixture of chain lengths derived from coconut or palm kernel fat or is derived synthetically by olefin build up or OXO alcohols synthesis.

Preferred groups for  $\text{R}_2\text{R}_3$  and  $\text{R}_4$  are methyl and hydroxyethyl groups and the anion  $X$  may be selected from  
 30 halide, methosulphate, acetate and phosphate ions.

Examples of suitable quaternary ammonium compounds of formulae (i) for use herein are:

- coconut trimethyl ammonium chloride or bromide;
- coconut methyl dihydroxyethyl ammonium chloride or bromide;
- 35 decyl triethyl ammonium chloride;
- decyl dimethyl hydroxyethyl ammonium chloride or bromide;
- $\text{C}_{12}$ -15 dimethyl hydroxyethyl ammonium chloride or bromide;

coconut dimethyl hydroxyethyl ammonium chloride or bromide;  
myristyl trimethyl ammonium methyl sulphate;  
lauryl dimethyl benzyl ammonium chloride or bromide;  
lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium chloride or bromide;  
5 choline esters (compounds of formula (i) wherein R<sub>1</sub> is

CH<sub>2</sub>-CH<sub>2</sub>-O-C-C<sub>12-14</sub> alkyl and R<sub>2</sub>R<sub>3</sub>R<sub>4</sub> are methyl).



10

di-alkyl imidazolines [compounds of formula (i)].

Other cationic surfactants useful herein are also described in US 4,228,044 and in EP 000 224.

When included therein, the laundry detergent compositions  
15 of the present invention typically comprise from 0.2% to about 25%, preferably from about 1% to about 8% by weight of such cationic surfactants.

Ampholytic surfactants are also suitable for use in the laundry detergent compositions of the present invention. These  
20 surfactants can be broadly described as aliphatic derivatives of secondary or tertiary amines, or aliphatic derivatives of heterocyclic secondary and tertiary amines in which the aliphatic radical can be straight- or branched-chain. One of the aliphatic substituents contains at least about 8 carbon  
25 atoms, typically from about 8 to about 18 carbon atoms, and at least one contains an anionic water-solubilizing group, e.g. carboxy, sulfonate, sulfate. See US 3,929,678 (column 19, lines 18-35) for examples of ampholytic surfactants.

When included therein, the laundry detergent compositions  
30 of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such ampholytic surfactants.

Zwitterionic surfactants are also suitable for use in laundry detergent compositions. These surfactants can be  
35 broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary

phosphonium or tertiary sulfonium compounds. See US 3,929,678 (column 19, line 38 through column 22, line 48) for examples of zwitterionic surfactants.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such zwitterionic surfactants.

Semi-polar nonionic surfactants are a special category of nonionic surfactants which include water-soluble amine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; watersoluble phosphine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; and water-soluble sulfoxides containing one alkyl moiety from about 10 to about 18 carbon atoms and a moiety selected from the group consisting of alkyl and hydroxyalkyl moieties of from about 1 to about 3 carbon atoms.

Semi-polar nonionic detergent surfactants include the amine oxide surfactants having the formula:



wherein  $\text{R}^3$  is an alkyl, hydroxyalkyl, or alkyl phenyl group or mixtures thereof containing from about 8 to about 22 carbon atoms;  $\text{R}^4$  is an alkylene or hydroxyalkylene group containing from about 2 to about 3 carbon atoms or mixtures thereof;  $x$  is from 0 to about 3; and each  $\text{R}^5$  is an alkyl or hydroxyalkyl group containing from about 1 to about 3 carbon atoms or a polyethylene oxide group containing from about 1 to about 3 ethylene oxide groups. The  $\text{R}^5$  groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.

These amine oxide surfactants in particular include C<sub>10</sub>-C<sub>18</sub> alkyl dimethyl amine oxides and C<sub>8</sub>-C<sub>12</sub> alkoxy ethyl dihydroxy ethyl amine oxides.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such semi-polar nonionic surfactants.

#### Builder system

10 The compositions according to the present invention may further comprise a builder system. Any conventional builder system is suitable for use herein including aluminosilicate materials, silicates, polycarboxylates and fatty acids, materials such as ethylenediamine tetraacetate, metal ion  
15 sequestrants such as aminopolyphosphonates, particularly ethylenediamine tetramethylene phosphonic acid and diethylene triamine pentamethylenephosphonic acid. Though less preferred for obvious environmental reasons, phosphate builders can also be used herein.

20 Suitable builders can be an inorganic ion exchange material, commonly an inorganic hydrated aluminosilicate material, more particularly a hydrated synthetic zeolite such as hydrated zeolite A, X, B, HS or MAP.

Another suitable inorganic builder material is layered  
25 silicate, e.g. SKS-6 (Hoechst). SKS-6 is a crystalline layered silicate consisting of sodium silicate (Na<sub>2</sub>Si<sub>2</sub>O<sub>5</sub>).

Suitable polycarboxylates containing one carboxy group include lactic acid, glycolic acid and ether derivatives thereof as disclosed in Belgian Patent Nos. 831,368, 821,369  
30 and 821,370. Polycarboxylates containing two carboxy groups include the water-soluble salts of succinic acid, malonic acid, (ethylenedioxy) diacetic acid, maleic acid, diglycollic acid, tartaric acid, tartronic acid and fumaric acid, as well as the ether carboxylates described in German Offenle-enschrift  
35 2,446,686, and 2,446,487, US 3,935,257 and the sulfinyl carboxylates described in Belgian Patent No. 840,623.

Polycarboxylates containing three carboxy groups include, in particular, water-soluble citrates, aconitrates and



citraconates as well as succinate derivatives such as the carboxymethyloxysuccinates described in British Patent No. 1,379,241, lactoxysuccinates described in Netherlands Application 7205873, and the oxypolycarboxylate materials such  
5 as 2-oxa-1,1,3-propane tricarboxylates described in British Patent No. 1,387,447.

Polycarboxylates containing four carboxy groups include oxydisuccinates disclosed in British Patent No. 1,261,829, 1,1,2,2,-ethane tetracarboxylates, 1,1,3,3-propane  
10 tetracarboxylates containing sulfo substituents include the sulfosuccinate derivatives disclosed in British Patent Nos. 1,398,421 and 1,398,422 and in US 3,936,448, and the sulfonated pyrolysed citrates described in British Patent No. 1,082,179, while polycarboxylates containing phosphone substituents are  
15 disclosed in British Patent No. 1,439,000.

Alicyclic and heterocyclic polycarboxylates include cyclopentane-cis,cis-cis-tetracarboxylates, cyclopentadienide pentacarboxylates, 2,3,4,5-tetrahydro-furan - cis, cis, cis-tetracarboxylates, 2,5-tetrahydro-furan-cis, discarboxylates,  
20 2,2,5,5,-tetrahydrofuran - tetracarboxylates, 1,2,3,4,5,6-hexane - hexacarboxylates and carboxymethyl derivatives of polyhydric alcohols such as sorbitol, mannitol and xylitol. Aromatic polycarboxylates include mellitic acid, pyromellitic acid and the phthalic acid derivatives disclosed in British  
25 Patent No. 1,425,343.

Of the above, the preferred polycarboxylates are hydroxycarboxylates containing up to three carboxy groups per molecule, more particularly citrates.

Preferred builder systems for use in the present  
30 compositions include a mixture of a water-insoluble aluminosilicate builder such as zeolite A or of a layered silicate (SKS-6), and a water-soluble carboxylate chelating agent such as citric acid.

A suitable chelant for inclusion in the detergent  
35 compositions in accordance with the invention is ethylenediamine-N,N'-disuccinic acid (EDDS) or the alkali metal, alkaline earth metal, ammonium, or substituted ammonium salts thereof, or mixtures thereof. Preferred EDDS compounds

are the free acid form and the sodium or magnesium salt thereof. Examples of such preferred sodium salts of EDDS include  $\text{Na}_2\text{EDDS}$  and  $\text{Na}_4\text{EDDS}$ . Examples of such preferred magnesium salts of EDDS include  $\text{MgEDDS}$  and  $\text{Mg}_2\text{EDDS}$ . The  
5 magnesium salts are the most preferred for inclusion in compositions in accordance with the invention.

Preferred builder systems include a mixture of a water-insoluble aluminosilicate builder such as zeolite A, and a water soluble carboxylate chelating agent such as citric acid.

10 Other builder materials that can form part of the builder system for use in granular compositions include inorganic materials such as alkali metal carbonates, bicarbonates, silicates, and organic materials such as the organic phosphonates, amino polyalkylene phosphonates and amino  
15 polycarboxylates.

Other suitable water-soluble organic salts are the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms.

20 Polymers of this type are disclosed in GB-A-1,596,756. Examples of such salts are polyacrylates of MW 2000-5000 and their copolymers with maleic anhydride, such copolymers having a molecular weight of from 20,000 to 70,000, especially about 40,000.

25 Detergency builder salts are normally included in amounts of from 5% to 80% by weight of the composition. Preferred levels of builder for liquid detergents are from 5% to 30%.

### Enzymes

30 Preferred detergent compositions, in addition to the enzyme preparation of the invention, comprise other enzyme(s) which provides cleaning performance and/or fabric care benefits.

Such enzymes include other proteases, lipases, cutinases,  
35 amylases, cellulases, peroxidases, oxidases (e.g. laccases).

Proteases: Any other protease suitable for use in alkaline solutions can be used. Suitable proteases include those of animal,

vegetable or microbial origin. Microbial origin is preferred. Chemically or genetically modified mutants are included. The protease may be a serine protease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of  
5 alkaline proteases are subtilisins, especially those derived from Bacillus, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168 (described in WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the Fusarium protease  
10 described in WO 89/06270.

Preferred commercially available protease enzymes include those sold under the trade names Alcalase, Savinase, Primase, Durazym, and Esperase by Novo Nordisk A/S (Denmark), those sold under the tradename Maxatase, Maxacal, Maxapem, Properase,  
15 Purafect and Purafect OXP by Genencor International, and those sold under the tradename Opticlean and Optimase by Solvay Enzymes. Protease enzymes may be incorporated into the compositions in accordance with the invention at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein  
20 by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

25

Lipases: Any lipase suitable for use in alkaline solutions can be used. Suitable lipases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included.

30 Examples of useful lipases include a Humicola lanuginosa lipase, e.g., as described in EP 258 068 and EP 305 216, a Rhizomucor miehei lipase, e.g., as described in EP 238 023, a Candida lipase, such as a C. antarctica lipase, e.g., the C. antarctica lipase A or B described in EP 214 761, a Pseudomonas  
35 lipase such as a P. alcaligenes and P. pseudoalcaligenes lipase, e.g., as described in EP 218 272, a P. cepacia lipase, e.g., as described in EP 331 376, a P. stutzeri lipase, e.g., as disclosed in GB 1,372,034, a P. fluorescens lipase, a Bacil-

lus lipase, e.g., a B. subtilis lipase (Dartois et al., (1993), Biochemica et Biophysica acta 1131, 253-260), a B. stearo-thermophilus lipase (JP 64/744992) and a B. pumilus lipase (WO 91/16422).

5 Furthermore, a number of cloned lipases may be useful, including the Penicillium camembertii lipase described by Yamaguchi et al., (1991), Gene 103, 61-67), the Geotricum candidum lipase (Schimada, Y. et al., (1989), J. Biochem., 106, 383-388), and various Rhizopus lipases such as a R. delemar  
10 lipase (Hass, M.J et al., (1991), Gene 109, 117-113), a R. niveus lipase (Kugimiya et al., (1992), Biosci. Biotech. Biochem. 56, 716-719) and a R. oryzae lipase.

Other types of lipolytic enzymes such as cutinases may also be useful, e.g., a cutinase derived from Pseudomonas  
15 mendocina as described in WO 88/09367, or a cutinase derived from Fusarium solani pisi (e.g. described in WO 90/09446).

Especially suitable lipases are lipases such as M1 Lipase<sup>TM</sup>, Luma fast<sup>TM</sup> and Lipomax<sup>TM</sup> (Genencor), Lipolase<sup>TM</sup> and Lipolase Ultra<sup>TM</sup> (Novo Nordisk A/S), and Lipase P "Amano"  
20 (Amano Pharmaceutical Co. Ltd.).

The lipases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition,  
25 more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

30 Amylases: Any amylase ( $\alpha$  and/or  $\beta$ ) suitable for use in alkaline solutions can be used. Suitable amylases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. Amylases include, for example,  $\alpha$ -amylases obtained from a special strain of B. licheniformis, described  
35 in more detail in GB 1,296,839. Commercially available amylases are Duramyl<sup>TM</sup>, Termamyl<sup>TM</sup>, Fungamyl<sup>TM</sup> and BAN<sup>TM</sup> (available from

Novo Nordisk A/S) and Rapidase<sup>TM</sup> and Maxamyl P<sup>TM</sup> (available from Genencor).

The amylases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Cellulases: Any cellulase suitable for use in alkaline solutions can be used. Suitable cellulases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. Suitable cellulases are disclosed in US 4,435,307, which discloses fungal cellulases produced from Humicola insolens. Especially suitable cellulases are the cellulases having colour care benefits. Examples of such cellulases are cellulases described in European patent application No. 0 495 257.

Commercially available cellulases include Celluzyme<sup>TM</sup> produced by a strain of Humicola insolens, (Novo Nordisk A/S), and KAC-500(B)<sup>TM</sup> (Kao Corporation).

Cellulases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Peroxidases/Oxidases: Peroxidase enzymes are used in combination with hydrogen peroxide or a source thereof (e.g. a percarbonate, perborate or persulfate). Oxidase enzymes are used in combination with oxygen. Both types of enzymes are used for "solution bleaching", i.e. to prevent transfer of a textile

dye from a dyed fabric to another fabric when said fabrics are washed together in a wash liquor, preferably together with an enhancing agent as described in e.g. WO 94/12621 and WO 95/01426. Suitable peroxidases/oxidases include those of plant, 5 bacterial or fungal origin. Chemically or genetically modified mutants are included.

Peroxidase and/or oxidase enzymes are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, 10 preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

15 Mixtures of the above mentioned enzymes are encompassed herein, in particular a mixture of a protease, an amylase, a lipase and/or a cellulase.

The enzyme of the invention, or any other enzyme incorporated in the detergent composition, is normally 20 incorporated in the detergent composition at a level from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level from 0.001% to 0.5% of enzyme protein by weight of the composition, 25 even more preferably at a level from 0.01% to 0.2% of enzyme protein by weight of the composition.

Bleaching agents: Additional optional detergent ingredients that can be included in the detergent compositions of the 30 present invention include bleaching agents such as PB1, PB4 and percarbonate with a particle size of 400-800 microns. These bleaching agent components can include one or more oxygen bleaching agents and, depending upon the bleaching agent chosen, one or more bleach activators. When present oxygen 35 bleaching compounds will typically be present at levels of from about 1% to about 25%. In general, bleaching compounds are optional added components in non-liquid formulations, e.g. granular detergents.

The bleaching agent component for use herein can be any of the bleaching agents useful for detergent compositions including oxygen bleaches as well as others known in the art.

The bleaching agent suitable for the present invention  
5 can be an activated or non-activated bleaching agent.

One category of oxygen bleaching agent that can be used encompasses percarboxylic acid bleaching agents and salts thereof. Suitable examples of this class of agents include magnesium monoperoxyphthalate hexahydrate, the magnesium salt  
10 of meta-chloro perbenzoic acid, 4-nonylamino-4-oxoperoxybutyric acid and diperoxydodecanedioic acid. Such bleaching agents are disclosed in US 4,483,781, US 740,446, EP 0 133 354 and US 4,412,934. Highly preferred bleaching agents also include 6-nonylamino-6-oxoperoxycaproic acid as described in US  
15 4,634,551.

Another category of bleaching agents that can be used encompasses the halogen bleaching agents. Examples of hypohalite bleaching agents, for example, include trichloro isocyanuric acid and the sodium and potassium  
20 dichloroisocyanurates and N-chloro and N-bromo alkane sulphonamides. Such materials are normally added at 0.5-10% by weight of the finished product, preferably 1-5% by weight.

The hydrogen peroxide releasing agents can be used in combination with bleach activators such as tetra-  
25 acetylenediamine (TAED), nonanoyloxybenzenesulfonate (NOBS, described in US 4,412,934), 3,5-trimethylhexanoyloxybenzenesulfonate (ISONOBS, described in EP 120 591) or pentaacetylglucose (PAG), which are perhydrolyzed to form a peracid as the active bleaching species, leading to improved  
30 bleaching effect. In addition, very suitable are the bleach activators C8(6-octanamido-caproyl) oxybenzenesulfonate, C9(6-nonanamido caproyl) oxybenzenesulfonate and C10 (6-decanamido caproyl) oxybenzenesulfonate or mixtures thereof. Also suitable activators are acylated citrate esters such as disclosed in  
35 European Patent Application No. 91870207.7.

Useful bleaching agents, including peroxyacids and bleaching systems comprising bleach activators and peroxygen

bleaching compounds for use in cleaning compositions according to the invention are described in application USSN 08/136,626.

The hydrogen peroxide may also be present by adding an enzymatic system (i.e. an enzyme and a substrate therefore) which is capable of generation of hydrogen peroxide at the beginning or during the washing and/or rinsing process. Such enzymatic systems are disclosed in European Patent Application EP 0 537 381.

Bleaching agents other than oxygen bleaching agents are also known in the art and can be utilized herein. One type of non-oxygen bleaching agent of particular interest includes photoactivated bleaching agents such as the sulfonated zinc and/or aluminium phthalocyanines. These materials can be deposited upon the substrate during the washing process. Upon irradiation with light, in the presence of oxygen, such as by hanging clothes out to dry in the daylight, the sulfonated zinc phthalocyanine is activated and, consequently, the substrate is bleached. Preferred zinc phthalocyanine and a photoactivated bleaching process are described in US 4,033,718. Typically, detergent composition will contain about 0.025% to about 1.25%, by weight, of sulfonated zinc phthalocyanine.

Bleaching agents may also comprise a manganese catalyst. The manganese catalyst may, e.g., be one of the compounds described in "Efficient manganese catalysts for low-temperature bleaching", Nature 369, 1994, pp. 637-639.

Suds suppressors: Another optional ingredient is a suds suppressor, exemplified by silicones, and silica-silicone mixtures. Silicones can generally be represented by alkylated polysiloxane materials, while silica is normally used in finely divided forms exemplified by silica aerogels and xerogels and hydrophobic silicas of various types. These materials can be incorporated as particulates, in which the suds suppressor is advantageously releasably incorporated in a water-soluble or waterdispersible, substantially non surface-active detergent impermeable carrier. Alternatively the suds suppressor can be dissolved or dispersed in a liquid carrier and applied by spraying on to one or more of the other components.



A preferred silicone suds controlling agent is disclosed in US 3,933,672. Other particularly useful suds suppressors are the self-emulsifying silicone suds suppressors, described in German Patent Application DTOS 2,646,126. An example of such a compound is DC-544, commercially available from Dow Corning, which is a siloxane-glycol copolymer. Especially preferred suds controlling agent are the suds suppressor system comprising a mixture of silicone oils and 2-alkyl-alkanols. Suitable 2-alkyl-alkanols are 2-butyl-octanol which are commercially available under the trade name Isofol 12 R.

Such suds suppressor system are described in European Patent Application EP 0 593 841.

Especially preferred silicone suds controlling agents are described in European Patent Application No. 92201649.8. Said compositions can comprise a silicone/ silica mixture in combination with fumed nonporous silica such as Aerosil<sup>R</sup>.

The suds suppressors described above are normally employed at levels of from 0.001% to 2% by weight of the composition, preferably from 0.01% to 1% by weight.

20

Other components: Other components used in detergent compositions may be employed such as soil-suspending agents, soil-releasing agents, optical brighteners, abrasives, bactericides, tarnish inhibitors, coloring agents, and/or encapsulated or nonencapsulated perfumes.

Especially suitable encapsulating materials are water soluble capsules which consist of a matrix of polysaccharide and polyhydroxy compounds such as described in GB 1,464,616.


Other suitable water soluble encapsulating materials comprise dextrans derived from ungelatinized starch acid esters of substituted dicarboxylic acids such as described in US 3,455,838. These acid-ester dextrans are, preferably, prepared from such starches as waxy maize, waxy sorghum, sago, tapioca and potato. Suitable examples of said encapsulation materials include N-Lok manufactured by National Starch. The N-Lok encapsulating material consists of a modified maize starch and glucose. The starch is modified by adding monofunctional substituted groups such as octenyl succinic acid anhydride.

Antiredeposition and soil suspension agents suitable herein include cellulose derivatives such as methylcellulose, carboxymethylcellulose and hydroxyethylcellulose, and homo- or co-polymeric polycarboxylic acids or their salts. Polymers of 5 this type include the polyacrylates and maleic anhydride-acrylic acid copolymers previously mentioned as builders, as well as copolymers of maleic anhydride with ethylene, methylvinyl ether or methacrylic acid, the maleic anhydride constituting at least 20 mole percent of the copolymer. These 10 materials are normally used at levels of from 0.5% to 10% by weight, more preferably from 0.75% to 8%, most preferably from 1% to 6% by weight of the composition.

Preferred optical brighteners are anionic in character, examples of which are disodium 4,4'-bis-(2-diethanolamino-4- 15 anilino -s- triazin-6-ylamino)stilbene-2:2' disulphonate, disodium 4, - 4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino-stilbene-2:2' - disulphonate, disodium 4,4' - bis-(2,4-dianilino-s-triazin-6-ylamino)stilbene-2:2' - disulphonate, monosodium 4',4'' - bis-(2,4-dianilino-s-tri-azin-6 20 ylamino)stilbene-2-sulphonate, disodium 4,4' -bis-(2-anilino-4-(N-methyl-N-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2' - disulphonate, di-sodium 4,4' -bis-(4-phenyl-2,1,3-triazol-2-yl)-stilbene-2,2' disulphonate, di-so-dium 4,4'bis(2-anilino-4-(1-methyl-2-hydroxyethylamino)-s-triazin-6-ylami- 25 no)stilbene-2,2'disulphonate, sodium 2(stilbyl-4''-(naphtho-1',2':4,5)-1,2,3, - triazole-2''-sulphonate and 4,4'-bis(2-sulphostyryl)biphenyl.

Other useful polymeric materials are the polyethylene glycols, particularly those of molecular weight 1000-10000, 30 more particularly 2000 to 8000 and most preferably about 4000. These are used at levels of from 0.20% to 5% more preferably from 0.25% to 2.5% by weight. These polymers and the previously mentioned homo- or co-polymeric poly-carboxylate salts are valuable for improving whiteness maintenance, fabric ash 35 deposition, and cleaning performance on clay, proteinaceous and oxidizable soils in the presence of transition metal impurities.

Soil release agents useful in compositions of the present invention are conventionally copolymers or terpolymers of terephthalic acid with ethylene glycol and/or propylene glycol units in various arrangements. Examples of such polymers are disclosed in US 4,116,885 and 4,711,730 and EP 0 272 033. A particular preferred polymer in accordance with EP 0 272 033 has the formula:



where PEG is  $-(\text{OC}_2\text{H}_4)_n-$ , PO is  $(\text{OC}_3\text{H}_6\text{O})$  and T is  $(\text{pOOC}_6\text{H}_4\text{CO})$ .

Also very useful are modified polyesters as random copolymers of dimethyl terephthalate, dimethyl sulfoisophthalate, ethylene glycol and 1,2-propanediol, the end groups consisting primarily of sulphobenzoate and secondarily of mono esters of ethylene glycol and/or 1,2-propanediol. The target is to obtain a polymer capped at both end by sulphobenzoate groups, "primarily", in the present context most of said copolymers herein will be endcapped by sulphobenzoate groups. However, some copolymers will be less than fully capped, and therefore their end groups may consist of monoester of ethylene glycol and/or 1,2-propanediol, thereof consist "secondarily" of such species.

The selected polyesters herein contain about 46% by weight of dimethyl terephthalic acid, about 16% by weight of 1,2-propanediol, about 10% by weight ethylene glycol, about 13% by weight of dimethyl sulfo benzoic acid and about 15% by weight of sulfoisophthalic acid, and have a molecular weight of about 3.000. The polyesters and their method of preparation are described in detail in EP 311 342.

Softening agents: Fabric softening agents can also be incorporated into laundry detergent compositions in accordance with the present invention. These agents may be inorganic or organic in type. Inorganic softening agents are exemplified by the smectite clays disclosed in GB-A-1 400898 and in US

5,019,292. Organic fabric softening agents include the water insoluble tertiary amines as disclosed in GB-A1 514 276 and EP 0 011 340 and their combination with mono C<sub>12</sub>-C<sub>14</sub> quaternary ammonium salts are disclosed in EP-B-0 026 528 and di-long-chain amides as disclosed in EP 0 242 919. Other useful organic ingredients of fabric softening systems include high molecular weight polyethylene oxide materials as disclosed in EP 0 299 575 and 0 313 146.

Levels of smectite clay are normally in the range from 5% to 15%, more preferably from 8% to 12% by weight, with the material being added as a dry mixed component to the remainder of the formulation. Organic fabric softening agents such as the water-insoluble tertiary amines or dilong chain amide materials are incorporated at levels of from 0.5% to 5% by weight, normally from 1% to 3% by weight whilst the high molecular weight polyethylene oxide materials and the water soluble cationic materials are added at levels of from 0.1% to 2%, normally from 0.15% to 1.5% by weight. These materials are normally added to the spray dried portion of the composition, although in some instances it may be more convenient to add them as a dry mixed particulate, or spray them as molten liquid on to other solid components of the composition.

Polymeric dye-transfer inhibiting agents: The detergent compositions according to the present invention may also comprise from 0.001% to 10%, preferably from 0.01% to 2%, more preferably from 0.05% to 1% by weight of polymeric dye-transfer inhibiting agents. Said polymeric dye-transfer inhibiting agents are normally incorporated into detergent compositions in order to inhibit the transfer of dyes from colored fabrics onto fabrics washed therewith. These polymers have the ability of complexing or adsorbing the fugitive dyes washed out of dyed fabrics before the dyes have the opportunity to become attached to other articles in the wash.

Especially suitable polymeric dye-transfer inhibiting agents are polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylpyrrolidone

polymers, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof.

Addition of such polymers also enhances the performance of the enzymes according the invention.

5 The detergent composition according to the invention can be in liquid, paste, gels, bars or granular forms.

Non-dusting granulates may be produced, e.g., as disclosed in US 4,106,991 and 4,661,452 (both to Novo Industri A/S) and may optionally be coated by methods known in the art.  
10 Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molecular weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there  
15 are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591.

Granular compositions according to the present invention  
20 can also be in "compact form", i.e. they may have a relatively higher density than conventional granular detergents, i.e. from 550 to 950 g/l; in such case, the granular detergent compositions according to the present invention will contain a lower amount of "Inorganic filler salt", compared to  
25 conventional granular detergents; typical filler salts are alkaline earth metal salts of sulphates and chlorides, typically sodium sulphate; "Compact" detergent typically comprise not more than 10% filler salt. The liquid compositions according to the present invention can also be in "concentrated  
30 form", in such case, the liquid detergent compositions according to the present invention will contain a lower amount of water, compared to conventional liquid detergents. Typically, the water content of the concentrated liquid detergent is less than 30%, more preferably less than 20%, most preferably less  
35 than 10% by weight of the detergent compositions.

The compositions of the invention may for example, be formulated as hand and machine laundry detergent compositions including laundry additive compositions and compositions

suitable for use in the pretreatment of stained fabrics, rinse added fabric softener compositions, and compositions for use in general household hard surface cleaning operations and dishwashing operations.

5       The following examples are meant to exemplify compositions for the present invention, but are not necessarily meant to limit or otherwise define the scope of the invention.

      In the detergent compositions, the abbreviated component identifications have the following meanings:

10

LAS:           Sodium linear C<sub>12</sub> alkyl benzene sulphonate

TAS:           Sodium tallow alkyl sulphate

15 XYAS:        Sodium C<sub>1X</sub> - C<sub>1Y</sub> alkyl sulfate

SS:            Secondary soap surfactant of formula 2-butyl octanoic acid

20 25EY:        A C<sub>12</sub> - C<sub>15</sub> predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide

45EY:        A C<sub>14</sub> - C<sub>15</sub> predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide  
25

XYEYS:       C<sub>1X</sub> - C<sub>1Y</sub> sodium alkyl sulfate condensed with an average of Z moles of ethylene oxide per mole

30

Nonionic:     C<sub>13</sub> - C<sub>15</sub> mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5 sold under the tradename Plurafax LF404 by BASF  
35               GmbH

CFAA:         C<sub>12</sub> - C<sub>14</sub> alkyl N-methyl glucamide

- TFAA:  $C_{16} - C_{18}$  alkyl N-methyl glucamide
- Silicate: Amorphous Sodium Silicate ( $SiO_2:Na_2O$  ratio = 2.0)
- 5 NaSKS-6: Crystalline layered silicate of formula  $\delta-Na_2Si_2O_5$
- Carbonate: Anhydrous sodium carbonate
- 10 Phosphate: Sodium tripolyphosphate
- MA/AA: Copolymer of 1:4 maleic/acrylic acid, average molecular weight about 80,000
- 15 Polyacrylate: Polyacrylate homopolymer with an average molecular weight of 8,000 sold under the tradename PA30 by BASF GmbH
- 20 Zeolite A: Hydrated Sodium Aluminosilicate of formula  $Na_{12}(AlO_2SiO_2)_{12} \cdot 27H_2O$  having a primary particle size in the range from 1 to 10 micrometers
- Citrate: Tri-sodium citrate dihydrate
- 25 Citric: Citric Acid
- Perborate: Anhydrous sodium perborate monohydrate bleach, empirical formula  $NaBO_2 \cdot H_2O_2$
- 30 PB4: Anhydrous sodium perborate tetrahydrate
- Percarbonate: Anhydrous sodium percarbonate bleach of empirical formula  $2Na_2CO_3 \cdot 3H_2O_2$
- 35 TAED: Tetraacetyl ethylene diamine

- CMC: Sodium carboxymethyl cellulose
- 5 DETPMP: Diethylene triamine penta (methylene phosphonic acid), marketed by Monsanto under the Tradename Dequest 2060
- PVP: Polyvinylpyrrolidone polymer
- 10 EDDS: Ethylenediamine-N, N'-disuccinic acid, [S,S] isomer in the form of the sodium salt
- Suds 25% paraffin wax Mpt 50°C, 17% hydrophobic silica, 58% Suppressor: paraffin oil
- 15 Granular
- Suds: 12% Silicone/silica, 18% stearyl alcohol, 70% suppressor: starch in granular form
- 20 Sulphate: Anhydrous sodium sulphate
- HMWPEO: High molecular weight polyethylene oxide
- TAE 25: Tallow alcohol ethoxylate (25)

25

Detergent Example I

A granular fabric cleaning composition in accordance with the invention may be prepared as follows:

30	Sodium linear C <sub>12</sub> alkyl	6.5
	benzene sulfonate	
	Sodium sulfate	15.0
	Zeolite A	26.0
35	Sodium nitrilotriacetate	5.0



47

	Enzyme of the invention	0.1
	PVP	0.5
5	TAED	3.0
	Boric acid	4.0
	Perborate	18.0
10	Phenol sulphonate	0.1
	Minors	Up to 100

15 Detergent Example II

A compact granular fabric cleaning composition (density 800 g/l) in accord with the invention may be prepared as follows:

20	45AS	8.0
	25E3S	2.0
	25E5	3.0
	25E3	3.0
	TFAA	2.5
25	Zeolite A	17.0
	NaSKS-6	12.0
	Citric acid	3.0
	Carbonate	7.0
	MA/AA	5.0
30	CMC	0.4
	Enzyme of the invention	0.1
	TAED	6.0
	Percarbonate	22.0
	EDDS	0.3
35	Granular suds suppressor	3.5
	water/minors	Up to 100%

Detergent Example III

Granular fabric cleaning compositions in accordance with the invention which are especially useful in the laundering of coloured fabrics were prepared as follows:

5	LAS	10.7	-
	TAS	2.4	-
	TFAA	-	4.0
	45AS	3.1	10.0
	45E7	4.0	-
10	25E3S	-	3.0
	68E11	1.8	-
	25E5	-	8.0
	Citrate	15.0	7.0
	Carbonate	-	10
15	Citric acid	2.5	3.0
	Zeolite A	32.1	25.0
	Na-SKS-6	-	9.0
	MA/AA	5.0	5.0
	DETPMP	0.2	0.8
20	Enzyme of the invention	0.10	0.05
	Silicate	2.5	-
	Sulphate	5.2	3.0
	PVP	0.5	-
	Poly (4-vinylpyridine)-N-	-	0.2
25	Oxide/copolymer of vinyl-imidazole and vinyl-pyrrolidone		
	Perborate	1.0	-
30	Phenol sulfonate	0.2	-
	Water/Minors	Up to 100%	

Detergent Example IV

Granular fabric cleaning compositions in accordance with the invention which provide "Softening through the wash" capability may be prepared as follows:

35 the invention which provide "Softening through the wash" capability may be prepared as follows:

49

	45AS	-	10.0
	LAS	7.6	-
	68AS	1.3	-
	45E7	4.0	-
5	25E3	-	5.0
	Coco-alkyl-dimethyl hydroxy-ethyl ammonium chloride	1.4	1.0
	Citrate	5.0	3.0
10	Na-SKS-6	-	11.0
	Zeolite A	15.0	15.0
	MA/AA	4.0	4.0
	DETPMP	0.4	0.4
	Perborate	15.0	-
15	Percarbonate	-	15.0
	TAED	5.0	5.0
	Smectite clay	10.0	10.0
	HMWPEO	-	0.1
	Enzyme of the invention	0.10	0.05
20	Silicate	3.0	5.0
	Carbonate	10.0	10.0
	Granular suds suppressor	1.0	4.0
	CMC	0.2	0.1
	Water/Minors	Up to 100%	

25

Detergent Example V

Heavy duty liquid fabric cleaning compositions in accordance with the invention may be prepared as follows:

30		I	II
	LAS acid form	-	25.0
	Citric acid	5.0	2.0
	25AS acid form	8.0	-
	25AE2S acid form	3.0	-
35	25AE7	8.0	-
	CFAA	5	-
	DETPMP	1.0	1.0

50

	Fatty acid	8	-
	Oleic acid	-	1.0
	Ethanol	4.0	6.0
	Propanediol	2.0	6.0
5	Enzyme of the invention	0.10	0.05
	Coco-alkyl dimethyl	-	3.0
	hydroxy ethyl ammonium		
	chloride		
10	Smectite clay	-	5.0
	PVP	2.0	-
	Water / Minors	Up to 100%	

## LEATHER INDUSTRY APPLICATIONS

15 A subtilase of the invention may be used in the leather industry, in particular for use in depilation of skins.

In said application a subtilase variant of the invention is preferably used in an enzyme composition which further comprise another protease.

20 For a more detailed description of suitable other proteases see section relating to suitable enzymes for use in a detergent composition (*vide supra*).

## WOOL INDUSTRY APPLICATIONS

25 A subtilase of the invention may be used in the wool industry, in particular for use in cleaning of clothes comprising wool.

In said application a subtilase variant of the invention is preferably used in an enzyme composition which further comprise another protease.

30 For a more detailed description of suitable other proteases see section relating to suitable enzymes for use in a detergent composition (*vide supra*).

35 The invention is described in further detail in the following examples which are not in any way intended to limit the scope of the invention as claimed.

## MATERIALS AND METHODS

### Strains:

*B. subtilis* DN1885 (Diderichsen et al., 1990).

- 5        *B. lentus* 309 and 147 are specific strains of *Bacillus lentus*, deposited with the NCIB and accorded the accession numbers NCIB 10309 and 10147, and described in US Patent No. 3,723,250 incorporated by reference herein.

- 10        *E. coli* MC 1000 (M.J. Casadaban and S.N. Cohen (1980); *J. Mol. Biol.* **138** 179-207), was made  $r^-$ ,  $m^+$  by conventional methods and is also described in US Patent Application Serial No. 039,298.

### Plasmids:

- 15 pJS3: *E. coli* - *B. subtilis* shuttle vector containing a synthetic gene encoding for subtilase 309. (Described by Jacob Schiødt et al. in Protein and Peptide letters 3:39-44 (1996)).

pSX222: *B. subtilis* expression vector (Described in WO 96/34946).

20

### General molecular biology methods:

- Unless otherwise mentioned the DNA manipulations and transformations were performed using standard methods of molecular biology (Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995; Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for *Bacillus*". John Wiley and Sons, 1990).

- 30 Enzymes for DNA manipulations were used according to the specifications of the suppliers.

### Enzymes for DNA manipulations

- Unless otherwise mentioned all enzymes for DNA manipulations, such as e.g. restriction endonucleases, ligases etc., are obtained from New England Biolabs, Inc.

**Proteolytic Activity**

In the context of this invention proteolytic activity is expressed in Kilo NOVO Protease Units (KNPU). The activity is determined relatively to an enzyme standard (SAVINASE<sup>®</sup>), and the determination is based on the digestion of a dimethyl casein (DMC) solution by the proteolytic enzyme at standard conditions, i.e. 50°C, pH 8.3, 9 min. reaction time, 3 min. measuring time. A folder AF 220/1 is available upon request to Novo Nordisk A/S, Denmark, which folder is hereby included by reference.

A GU is a Glycine Unit, defined as the proteolytic enzyme activity which, under standard conditions, during a 15-minutes' incubation at 40 deg C, with N-acetyl casein as substrate, produces an amount of NH<sub>2</sub>-group equivalent to 1 mmole of glycine.

Enzyme activity can also be measured using the PNA assay, according to reaction with the soluble substrate succinyl-alanine-alanine-proline-phenyl-alanine-para-nitrophenol, which is described in the Journal of American Oil Chemists Society, Rothgeb, T.M., Goodlander, B.D., Garrison, P.H., and Smith, L.A., (1988).

**Fermentation:**

Fermentation of subtilase enzymes were performed at 30°C on a rotary shaking table (300 r.p.m.) in 500 ml baffled Erlenmeyer flasks containing 100 ml BPX medium for 5 days.

Consequently in order to make an e.g. 2 liter broth Erlenmeyer flasks were fermented simultaneously.

**Media:****BPX: Composition (per liter)**

	Potato starch	100g
	Ground barley	50g
	Soybean flour	20g
35	Na <sub>2</sub> HPO <sub>4</sub> X 12 H <sub>2</sub> O	9g
	Pluronic	0.1g
	Sodium caseinate	10g

The starch in the medium is liquified with  $\alpha$ -amylase and the medium is sterilized by heating at 120°C for 45 minutes. After sterilization the pH of the medium is adjusted to 9 by addition of NaHCO<sub>3</sub> to 0.1 M.

### EXAMPLES

#### **EXAMPLE 1**

10

#### Construction and Expression of Enzyme Variants:

##### *Site-directed mutagenesis:*

Subtilase 309 site-directed variants was made by the "Unique site elimination (USE)" or the "Uracil-USE" technique described respectively by Deng et al. (Anal. Biochem. 200:81-88 (1992)) and Markvardsen et al. (BioTechniques 18(3):371-372 (1995)).

The template plasmid was pJS3, or a analogue of this containing a variant of Subtilase 309, e.g. USE mutagenesis was performed on pJS3 analogue containing a gene encoding the G97E variant with a oligonucleotide directed to the construct of a A98V variant resulting in a final G97E+A98V Subtilase 309 variant.

The in pJS3 constructed Subtilase 309 variants was then subcloned into the *B.subtilis* pSX222 expression plasmid, using the restriction enzymes KpnI and MluI.

##### *Localized Random mutagenesis:*

The overall strategy to used to perform localized random mutagenesis was:

a mutagenic primer (oligonucleotide) was synthesized which corresponds to the part of the DNA sequence to be mutagenized except for the nucleotide(s) corresponding to amino acid codon(s) to be mutagenized.

Subsequently, the resulting mutagenic primer was used in a PCR reaction with a suitable opposite primer. The resulting PCR fragment was purified and digested and cloned into a *E.coli-B.subtilis* shuttle vector.

Alternatively and if necessary, the resulting PCR fragment is used in a second PCR reaction as a primer with a second suitable opposite primer so as to allow digestion and cloning of the mutagenized region into the shuttle vector. The  
5 PCR reactions are performed under normal conditions.

Following this strategy a localized random library was constructed in SAVINASE wherein positions V95, G97 and A98 was completely randomized.

One oligonucleotide was synthesized with 25% of each of  
10 the four bases (N) in the first and the second base at amino acid codons wanted to be mutagenized. The third nucleotide (the wobble base) in codons were synthesized with 50%G/50%C (S) to avoid two (TAA, TGA) of the three stop-codons .

The mutagenic primer (5'-GC GCT GAG CTA TAC GCT GTT AAA  
15 NNS CTA NNS NNS AGC GGT TCA GGT TCG GTC-3' (sense)) were used in a PCR reaction with a suitable anti-sense opposite primer, situated downstream of the MluI site in pJS3, and the plasmid pJS3 as template. This resulting PCR product was cloned into the pJS3 shuttle vector by using the restriction enzymes BlnI  
20 and MluI.

The in pJS3 constructed localized random library was then subcloned into the *B.subtilis* pSX222 expression plasmid, using the restriction enzymes KpnI and MluI.

The library prepared contained approximately 100,000 indi-  
25 vidual clones/library.

Ten randomly chosen colonies were sequenced to confirm the mutations designed.

In order to purify a subtilase variant of the invention the  
30 *B.subtilis* pSX222 expression plasmid comprising a variant of the invention was transformed into a competent *B. subtilis* strain and was fermented as described above in a medium containing 10 µg/ml Chloramphenicol (CAM).



**EXAMPLE 2**Purification of Enzyme Variants:

This procedure relates to purification of a 2 litre scale  
5 fermentation of the Subtilisin 147 enzyme, the Subtilisin 309  
enzyme or mutants thereof.

Approximately 1.6 litres of fermentation broth were  
centrifuged at 5000 rpm for 35 minutes in 1 litre beakers. The  
supernatants were adjusted to pH 6.5 using 10% acetic acid and  
10 filtered on Seitz Supra S100 filter plates.

The filtrates were concentrated to approximately 400 ml  
using an Amicon CH2A UF unit equipped with an Amicon S1Y10 UF  
cartridge. The UF concentrate was centrifuged and filtered  
prior to absorption at room temperature on a Bacitracin af-  
15 finity column at pH 7. The protease was eluted from the  
Bacitracin column at room temperature using 25% 2-propanol and  
1 M sodium chloride in a buffer solution with 0.01 dimethyl-  
glutaric acid, 0.1 M boric acid and 0.002 M calcium chloride  
adjusted to pH 7.

20 The fractions with protease activity from the Bacitracin  
purification step were combined and applied to a 750 ml  
Sephadex G25 column (5 cm dia.) equilibrated with a buffer  
containing 0.01 dimethylglutaric acid, 0.2 M boric acid and  
0.002 m calcium chloride adjusted to pH 6.5.

25 Fractions with proteolytic activity from the Sephadex  
G25 column were combined and applied to a 150 ml CM Sepharose  
CL 6B cation exchange column (5 cm dia.) equilibrated with a  
buffer containing 0.01 M dimethylglutaric acid, 0.2 M boric  
acid, and 0.002 M calcium chloride adjusted to pH 6.5.

30 The protease was eluted using a linear gradient of 0-0.1  
M sodium chloride in 2 litres of the same buffer (0-0.2 M  
sodium chloride in case of Subtilisin 147).

In a final purification step protease containing  
fractions from the CM Sepharose column were combined and con-  
35 centrated in an Amicon ultrafiltration cell equipped with a  
GR81PP membrane (from the Danish Sugar Factories Inc.).

By using the techniques of Example 1 for the construction and the above isolation procedure the following subtilisin 309 variants were produced and isolated:

V95T+G97T+A98E; and

5 G97E+A98V.

### EXAMPLE 3

#### 10 Wash Performance of Detergent Compositions Comprising Enzyme Variants

The following examples provide results from a number of washing tests that were conducted under the conditions indicated

#### EXPERIMENTAL CONDITIONS

15

Table VI: Experimental conditions for evaluation of Subtilisin 309 variants.

Detergent	Protease Model Detergent 95
Detergent dose	3.0 g/l
pH	10.5
Wash time	15 min.
Temperature	15°C
Water hardness	6°dH
Enzymes	Subtilisin 309 variants as listed below
Enzyme conc.	10 nM
Test system	150 ml glass beakers with a stirring rod
Textile/volume	5 textile pieces (Ø 2.5 cm) in 50 ml detergent

Test material	EMPA117 from Center for Testmaterials, Holland
---------------	---

The detergent used is a simple model formulation. pH is adjusted to 10.5 which is within the normal range for a powder detergent. The composition of model detergent 95 is as follows:

5

25% STP ( $\text{Na}_5\text{P}_3\text{O}_{10}$ )

25%  $\text{Na}_2\text{SO}_4$

10%  $\text{Na}_2\text{CO}_3$

20% LAS (Nansa 80S)

10 5.0% Nonionic tenside (Dobanol 25-7)

5.0%  $\text{Na}_2\text{Si}_2\text{O}_5$

0.5% Carboxymethylcellulose (CMC)

9.5% Water

15 Water hardness was adjusted by adding  $\text{CaCl}_2$  and  $\text{MgCl}_2$  ( $\text{Ca}^{2+}:\text{Mg}^{2+} = 2:1$ ) to deionized water (see also Surfactants in Consumer Products - Theory, Technology and Application, Springer Verlag 1986). pH of the detergent solution was adjusted to pH 10.5 by addition of HCl.

20

Measurement of reflectance (R) on the test material was done at 460 nm using a Macbeth ColorEye 7000 photometer (Macbeth, Division of Kollmorgen Instruments Corporation, Germany). The measurements were done according to the manufacturers protocol.

25

The wash performance of the Subtilisin 309 variants was evaluated by calculating a performance factor:

$$P = \frac{R_{\text{Variant}} - R_{\text{Blank}}}{R_{\text{Savinase}} - R_{\text{Blank}}}$$

30

P: Performance factor

$R_{\text{Variant}}$ : Reflectance of test material washed with variant

$R_{\text{Savinase}}$ : Reflectance of test material washed with Savinase®

R<sub>Blank</sub>: Reflectance of test material washed with no enzyme

The claimed Subtilisin 309 variants all have improved wash performance compared to Savinase® - i.e.  $P > 1$ .

5

The variants are divided into improvement classes designated with capital letters:

Class A:  $1 < P \leq 1.5$

10 Class B:  $1.5 < P \leq 2$

Class C:  $P > 2$

Table V: Subtilisin 309 variants and improvement classes.

Improvement class	Variants
A	V95T+G97T+A98E G97E+A98V
B	
C	

**PATENT CLAIMS**

1. A subtilase enzyme variant having improved wash performance in detergents, comprising modification(s) in position(s) 95, 97 and/or 98 (in BASBPN numbering).

2. A subtilase enzyme variant having improved wash performance in detergents, comprising at least one modification chosen from the group comprising (in BASBPN numbering):

10 95T+97T+98E

97E+98V; or

a variant comprising one or more conservative modification(s) in any of the above mentioned variants (e.g. a conservative modification of a 97E(*acidic a.a.*)+98V variant include a variant such as 97D(*acidic a.a.*)+98V.).

3. The subtilase enzyme variant according to claim 2, wherein the modification is chosen from the group comprising (in BASBPN numbering):

20 V95T+G97T+A98E

G97E+A98V; or

a variant comprising one or more conservative modification(s) in any of the above mentioned variants (e.g. a conservative modification of a G97E(*acidic a.a.*)+A98V variant include a variant such as G97D(*acidic a.a.*)+A98V.).

4. The variant of any of the claims 1 to 3, wherein the parent subtilase is chosen from the sub-group I-S1.

30 5. The variant of claim 4, wherein the parent subtilase is chosen from the group comprising ABSS168, BASBPN, BSSDY, and BLSCAR or functional variants thereof having retained the characteristic of sub-group I-S1.

35 6. The variant of any of the claims 1 to 3, wherein the parent subtilase is chosen from the sub-group I-S2.

7. The variant of claim 6, wherein the parent subtilase is chosen from the group comprising BLS147, BLS309, BAPB92, TVTHER AND BYSYAB or functional variants thereof having retained the characteristic of sub-group I-S2.

5

8. The variant of any of the claims above, wherein said modification(s) is/are combined with one or more modification(s) in any other position(s).

10 9. The variant of claim 8, wherein said modification(s) is/are combined with modification(s) in one or more of the positions 27, 36, 57, 76, 97, 101, 104, 120, 123, 167, 170, 206, 218, 222, 224, 235 and 274.

15 10. The variant of claim 9, wherein said subtilase belongs to the I-S2 sub-group and said further change is chosen from the group comprising K27R, \*36D, S57P, N76D, G97N, S101G, V104A, V104N, V104Y, H120D, N123S, Y167A, Y167I, R170S, R170L, R170N, Q206E, N218S, M222S, M222A, T224S, K235L, and T274A.

20

11. The variant of claim 10 comprising any of the variants V104N+S101G, K27R+V104Y+N123S+T274A, or N76D+V104A, or other combinations of these mutations (V104N, S101G, K27R, V104Y, N123S, T274A, N76D, V104A), in combination with any one or more  
25 of the substitutions, deletions and/or insertions mentioned in any of claims 1 to 10.

12. The subtilase variant of any of the preceding claims, wherein said modificaton(s) is/are combined with  
30 modification(s) in one or more of the positions 129, 131, 133 and 194.

13. The variant of claim 12, wherein said subtilase belongs to the I-S2 sub-group and said further change is chosen from the  
35 group comprising P129K, P131H, A133P, A133D and A194P.

14. An isolated DNA sequence encoding a subtilase variant of any of the claims 1 to 13.

15. An expression vector comprising an isolated DNA  
5 sequence of claim 14.

16. A microbial host cell transformed with an expression vector of claim 15.

10 17. The microbial host of claim 16, which is a bacterium, preferably a *Bacillus*, especially *B. lentus*.

18. The microbial host of claim 16, which is a fungus or yeast, preferably a filamentous fungus, especially an  
15 *Aspergillus*.

19. A method for producing a variant of any of claims 1 to 13, wherein a host of any of claims 16 to 18 is cultured under conditions conducive to the expression and secretion of said  
20 variant, and the variant is recovered.

20. A composition comprising a subtilase variant according to any of claims 1 to 13.

25 21. The composition according to claim 20, which additionally comprises a cellulase, lipase, cutinase, oxidoreductase, another protease, or an amylase.

22. The composition according to claim 20 or 21, wherein  
30 the composition is a detergent composition.

23. Use of a subtilase variant according to any of claims 1 to 13 or an enzyme composition according to any of claims 20 to 22 in a laundry and/or a dishwash detergent.

35

24. A process for the identification of a protease variant exhibiting improved wash performance in detergents, which



comprises effecting a mutation in DNA encoding a subtilase enzyme or its pre- or preproenzyme at one or more of the positions corresponding to amino acid (in BASBPN numbering):

V95T+G97T+A98E

5 G97E+A98V; or

a variant comprising one or more conservative modification(s) in any of the above mentioned variants;

transforming a Bacillus strain with said mutated DNA;

selecting strains producing such protease variants;

10 fermenting/growing such a strain;

recovering said protease variant, and

testing for improved wash performance in detergents.

## 1/2

## Alignment of subtilases

	1		10		20		23
{BASBPN}	.....AQ	SVP.....	YGVSQIKAPA	LH.SQGYTGS			
{BLS147}	.....Q	TVP.....	WGISFINTQQ	AH.NRGIFGN			
{BYSYAB}	.....Q	TVP.....	WGINRVQAPI	AQ.SRGFTGT			
{BAPB92}	.....AQ	SVP.....	WGISRVQAPA	AH.NRGLTGS			
{BSSDY}	.....AQ	TVP.....	YGIPLIKADK	VQ.AQGYKGA			
{TVTHER}	.....YTPNDPYFS	SRQ.....	YGPQKIQAPQ	AW.DIAE.GS			
{BLSAVI}	.....AQ	SVP.....	WGISRVQAPA	AH.NRGLTGS			
{BSISP1}	MNGEIRLIPY	VTNEQIMDVN	ELP.....	EGIKVIKAPK	MW.AKGVKVK		
{BSEPR}	.....SDGTDSDN	FEQ.....	WNLEPIQVKQ	AW.KAGLTGK			
{JP170}	LRGLEQIAQY	ATNNDVLYVT	PKPEYEV LND	VARGIVKADV	AQNNFGLYGO		
	30		40		50		60
	3234						
{BASBPN}	NVKVAVIDSG	IDSS.....	HPDLK..VAG	GASMVPSETN	...PFQDNNS		
{BLS147}	GARVAVLDTG	IAS.....	HPDLR..IAG	GASFISSEP.	...SYHDNNG		
{BYSYAB}	GVRVAVLDTG	ISN.....	HADLR..IRG	GASFPVGEF.	...NISDGNG		
{BAPB92}	GKVAVLDTG	IST.....	HPDLN..IRG	GASFPVGEF.	...STQDGNG		
{BSSDY}	NVKVGIIDTG	IAAS.....	HTDLK..VVG	GASFVSGES.	...YNTDGNG		
{TVTHER}	GAKIAIVDTG	VQSN.....	HPDLAGKVVG	GWDFVDNDS.	...TPQDNGG		
{BLSAVI}	GKVAVLDTG	IST.....	HPDLN..IRG	GASFPVGEF.	...STQDGNG		
{BSISP1}	NIKVAVLDTG	CDTS.....	HPDLKNQIIG	GKNFSDDDGG	KEDAI SDYNG		
{BSEPR}	NIKIAVIDSG	ISP.....	HDDLK..IAG	GYSVSVYS.	...SYKDDNG		
{JP170}	GQIVAVADTG	LDTGRNDSSM	HEAFRGKITA	LYALGRTNN.	....ANDPNG		
	70		80		90		100
	646566		83				110
{BASBPN}	HGTHVAGTVA	ALNN.SIGVL	GVAPSASLYA	VKVLG.ADGS	GQYSWIING.		
{BLS147}	HGTHVAGTIA	ALNN.SIGVL	GVRPSADLYA	LKVLD.RNGS	GSLASVAQG.		
{BYSYAB}	HGTQVAGTIA	ALNN.SIGVL	GVAPNVDLYG	VKVLG.ASGS	GSISGIAQG.		
{BAPB92}	HGTHVAGTIA	ALNN.SIGVL	GVAPNAELYA	VKVLG.ASGS	GSVSSIAQG.		
{BSSDY}	HGTHVAGTVA	ALDN.TTGVL	GVAPNVSLYA	IKVLN.SSGS	GTYSIVVSG.		
{TVTHER}	HGTHCAGIAA	AVTNNSTGIA	GTAPKASILA	VRVLD.NSGS	GTWTAVANG.		
{BLSAVI}	HGTHVAGTIA	ALNN.SIGVL	GVAPSAELYA	VKVLG.ASGS	GSVSSIAQG.		
{BSISP1}	HGTHVAGTIA	ANDS.NGGIA	GVAPASLLI	VKVLGGNGS	GQYEWIING.		
{BSEPR}	HGTHVAGIIG	AKHN.GYGID	GIAPEAQIYA	VKALD.QNGS	GDLQSLQGG.		
{JP170}	HGTHVAGSVL	GNAT..N..K	GMAPQANLVF	QSIMDSGGGL	GGLPANLQTL		
	120		130		140		150
		125127			146		154155
{BASBPN}	IEWAIANNMD	VINMSLGGPS	G..SAALKAA	VDKAVASG.V	VVAAAAGNEG		
{BLS147}	IEWAINNNMH	IINMSLGSTS	G..SSTLELA	VNRANNAG.I	LLVGAAGNTG		
{BYSYAB}	LQWAANNMGH	IANMSLGSSA	G..SATMEQA	VNQATASG.V	LVVAASGNNG		
{BAPB92}	LEWAGNNGMH	VANLSLGGSPS	P..SATLEQA	VNSATSRG.V	LVVAASGNNG		
{BSSDY}	IEWATQNGLD	VINMSLGGPS	G..STALKQA	VDKAYASG.I	VVAAAAGNSG		
{TVTHER}	ITYAADQGAK	VISLSLGGTV	G..NSGLQQA	VNYAWNKG.S	VVAAAAGNAG		
{BLSAVI}	LEWAGNNGMH	VANLSLGGSPS	P..SATLEQA	VNSATSRG.V	LVVAASGNNG		
{BSISP1}	INYAVEQKVD	IISMSLGGPS	D..VPELEEA	VKNAVKNV.V	LVVCAAGNEG		
{BSEPR}	IDWSIANRMD	IVNMSLGTTS	D..SKILHDA	VNKAYEQG.V	LLVAASGNNG		
{JP170}	FSQAYSAGAR	IHTNSWGAPV	NGAYTTDSRN	VDDYVRKNDM	TILFAAGNEG		

Fig. 1

2/2

	160	170	180	190
{BASBPN}	TSGS.SSTVG	YPGKYPSVIA	VGAVD.....	SSNQ RASFSSVG..
{BLS147}	RQG.....VN	YPARYSGVMA	VAVD.....	QNGQ RASFSTYG..
{BYSYAB}	AGN.....VG	FPARYANAMA	VGATD.....	QNNN RATFSQYG..
{BAPB92}	AGS.....IS	YPARYANAMA	VGATD.....	QNNN RASFQYG..
{BSSDY}	SSGS.QNTIG	YPAKYDSVIA	VGAVD.....	SNKN RASFSSVG..
{TVTHER}	NTAP.....N	YPAYYSNAIA	VASTD.....	QNDN KSSFSTYG..
{BLSAVI}	AGS.....IS	YPARYANAMA	VGATD.....	QNNN RASFQYG..
{BSISP1}	DGDERTEELS	YPAAYNEVIA	VGSVS.....	VARE LSEFSNAN..
{BSEPR}	NGKP....VN	YPAAYSSVVA	VSATN.....	EKNQ LASFSTTG..
{JP170}	PGSG...TIS	APGTAKNAIT	VGATENLRPS	FGSYADNINH VAQFSSRGPT

	200	210	220
			219 221 225
{BASBPN}	....PELDVM	APGVSIQSTL	PGNK..... YGAY NGTSMASPHV
{BLS147}	....PEIEIS	APGVNVNSTY	TGNR..... YVSL SGTSMATPHV
{BYSYAB}	....AGLDIV	APGVGVQSTV	PGNG..... YASF NGTSMATPHV
{BAPB92}	....AGLDIV	APGVNVQSTY	PGST..... YASL NGTSMATPHV
{BSSDY}	....AELEVVM	APGVSVYSTY	PSNT..... YTSL NGTSMASPHV
{TVTHER}	....SVVDVA	APGSKIYSTY	PTST..... YASL SGTSMATPHV
{BLSAVI}	....AGLDIV	APGVNVQSTY	PGST..... YASL NGTSMATPHV
{BSISP1}	....KEIDLV	APGENILSTL	PNKK..... YGKL TGTSMAPHV
{BSEPR}	....DEVEFS	APGTNITSTY	LNQY..... YATG SGTSQLATPHA
{JP170}	RDGRIKPDVM	APGTIILSAR	SSLAPDSSFW ANHDSKYAYM GGTSMATPIV

	230	240	250	260
{BASBPN}	AGAAALILSK	HP.....NWT	NTQVRSSLEN	TTTKLGDSF. ..YYGKGLIN
{BLS147}	AGVAALVKSR	YP.....SYT	NNQIRQRINQ	TATYLGSPS. ..LYGNGLVH
{BYSYAB}	AGVAALVKQK	NP.....SWS	NVQIRNHLKN	TATNLGNTT. ..QFGSGLVN
{BAPB92}	AGAAALVKQK	NP.....SWS	NVQIRNHLKN	TATSLGSTN. ..LYGSGLVN
{BSSDY}	AGAAALILSK	YP.....TLS	ASQVRNRLSS	TATNLGDSF. ..YYGKGLIN
{TVTHER}	AGVAGLLASQ	.....GRS	ASNIRAAIEN	TADKISGTG. .TYWAKGRVN
{BLSAVI}	AGAAALVKQK	NP.....SWS	NVQIRNHLKN	TATSLGSTN. ..LYGSGLVN
{BSISP1}	SGALALIKSY	EEESFQKLS	ESEVFAQLIR	RTLPLDIAKT ..LAGNGFLY
{BSEPR}	AAMFALLKQR	DP.....AET	NVQLREEMRK	NIVDLGTAGR DQQFGYGLIQ
{JP170}	AGNVAQLREH	FVKNRGVTPK	PSLLKAALIA	GAADVGLGFP NGNQGWGRVT

	270
{BASBPN}	VQAAAQ.
{BLS147}	AGRATQ.
{BYSYAB}	AEAATR.
{BAPB92}	AEAATR.
{BSSDY}	VEAAAQ.
{TVTHER}	AYKAVQY
{BLSAVI}	AEAATR.
{BSISP1}	LTAPDEL
{BSEPR}	YKAQATD
{JP170}	LDKSLNV

Fig. 1 (continued)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 98/00361

## A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C12N 9/54

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, CA, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9402618 A1 (GIST-BROCADES N.V.), 3 February 1994 (03.02.94) --	1-24
X	EP 0405901 A1 (UNILEVER PLC), 2 January 1991 (02.01.91), See claim 33 --	1-24
X	WO 9530011 A2 (THE PROCTER & GAMBLE COMPANY), 9 November 1995 (09.11.95), See claim 9B --	1-24
X	EP 0251446 A2 (GENENTECH, INC.), 7 January 1988 (07.01.88), See claim 3 -- -----	1-24

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

8 December 1998

10 -12- 1998

Name and mailing address of the ISA/  
Swedish Patent Office

Authorized officer

# INTERNATIONAL SEARCH REPORT

Information on patent family members

03/11/98

International application No.

PCT/DK 98/00361

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9402618 A1	03/02/94	AU 682314 B	02/10/97
		AU 4700693 A	14/02/94
		CA 2139928 A	03/02/94
		EP 0717778 A	26/06/96
		FI 950168 A	10/03/95
		JP 7508887 T	05/10/95
		PL 307132 A	02/05/95
EP 0405901 A1	02/01/91	CA 2034486 A	27/12/90
		JP 4500385 T	23/01/92
		US 5665587 A	09/09/97
		WO 9100334 A	10/01/91
		EP 0405902 A	02/01/91
		JP 4500384 T	23/01/92
		WO 9100335 A	10/01/91
WO 9530011 A2	09/11/95	AU 1253695 A	04/05/95
		AU 2294495 A	29/11/95
		AU 8079794 A	04/05/95
		BR 9407833 A	13/05/97
		BR 9407834 A	13/05/97
		BR 9507605 A	19/08/97
		CA 2173105 A	20/04/95
		CA 2173106 A	20/04/95
		CA 2189427 A	09/11/95
		CN 1137288 A	04/12/96
		CN 1137289 A	04/12/96
		CN 1151187 A	04/06/97
		CZ 9601052 A	11/09/96
		CZ 9601053 A	11/09/96
		CZ 9603217 A	14/05/97
		EP 0723579 A	31/07/96
		EP 0723580 A	31/07/96
		EP 0758389 A	19/02/97
		HU 75558 A	28/05/97
		HU 75984 A	28/05/97
		HU 76538 A	29/09/97
		HU 9600960 D	00/00/00
		HU 9600961 D	00/00/00
		HU 9603030 D	00/00/00
		JP 9503818 T	15/04/97
		JP 9505841 T	10/06/97
		JP 9512434 T	16/12/97
		US 5677272 A	14/10/97
		US 5679630 A	21/10/97
		WO 9510591 A	20/04/95
		WO 9510592 A	20/04/95

# INTERNATIONAL SEARCH REPORT

Information on patent family members

03/11/98

International application No.

PCT/DK 98/00361

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0251446 A2	07/01/88	SE 0251446 T3	
		AT 116365 T	15/01/95
		AU 614929 B	19/09/91
		AU 7228187 A	05/11/87
		DE 3750916 D,T	01/06/95
		DK 221787 A	31/10/87
		ES 2068181 T	16/04/95
		GR 3015388 T	30/06/95
		IE 65767 B	15/11/95
		JP 1085075 A	30/03/89
		JP 2647845 B	27/08/97
		JP 2772295 B	02/07/98
		JP 10042876 A	17/02/98
		NZ 240670 A	26/05/95
		SG 9590699 A	01/09/95
		US 5182204 A	26/01/93
		US 5185258 A	09/02/93
		US 5204015 A	20/04/93
		US 5371008 A	06/12/94
		US 5371190 A	06/12/94
		US 5472855 A	05/12/95
		US 5652136 A	29/07/97
		US 5700676 A	23/12/97
		US 5763257 A	09/06/98
		US 5801038 A	01/09/98
		US 5352594 A	04/10/94